

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification: C12N 15/12, C07K 14/47, A61K 38/17, C07K 16/18, C12Q 1/68, A61K 31/70	A2	(11) International Publication Number: WO 98/06841 (43) International Publication Date: 19 February 1998 (19.02.98)
(21) International Application Number: PCT/US97/13469 (22) International Filing Date: 24 July 1997 (24.07.97) (30) Priority Data: 08/700,607 12 August 1996 (12.08.96) US (60) Parent Application or Grant (63) Related by Continuation US 08/700,607 (CIP) Filed on 12 August 1996 (12.08.96) (71) Applicant (for all designated States except US): INCYTE PHARMACEUTICALS, INC. [US/US]; 3174 Porter Drive, Palo Alto, CA 94304 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): BANDMAN, Olga [US/US]; 366 Anna Avenue, Mountain View, CA 94043 (US). AU-YOUNG, Janice [US/US]; 1419 Kains Avenue, Berkeley, CA 94702 (US). GOLI, Surya, K. [IN/US]; 620 Iris Avenue #338, Sunnyvale, CA 94086 (US). HILLMAN, Jennifer [US/US]; 230 Monroe Drive #12, Mountain View, CA 94040 (US).		(74) Agent: BILLINGS, Lucy, J.; Incyte Pharmaceuticals, Inc., 3174 Porter Drive, Palo Alto, CA 94304 (US). (81) Designated States: AT, AU, BR, CA, CH, CN, DE, DK, ES, FI, GB, IL, JP, KR, MX, NO, NZ, RU, SE, SG, US, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: TWO HUMAN NSP-LIKE PROTEINS		
(57) Abstract <p>The present invention provides polynucleotides which identify and encode two human NSP-like proteins (NSPLP). The invention provides for genetically engineered expression vectors and host cells comprising the nucleic acid sequences encoding NSPLP. The invention also provides for the use of substantially purified NSPLP, antagonists, and in pharmaceutical compositions for the treatment of diseases associated with the expression of NSPLP. Additionally, the invention provides for the use of antisense molecules to NSPLP in pharmaceutical compositions for treatment of diseases associated with the expression of NSPLP. The invention also describes diagnostic assays which utilize diagnostic compositions comprising the polynucleotide, fragments or the complement thereof, which hybridize with the genomic sequence or the transcript of polynucleotides encoding NSPLP or anti-NSPLP antibodies which specifically bind to NSPLP.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

TWO HUMAN NSP-LIKE PROTEINS

5 The present invention relates to nucleic acid and amino acid sequences of two novel human NSP-like proteins and to the use of these sequences in the diagnosis, study, prevention and treatment of disease.

BACKGROUND ART

10 Neuroendocrine-specific proteins (NSP-A, NSP-B, and NSP-C) are a recently characterized group of membrane-anchored endoplasmic reticulum (ER) proteins that share identical carboxy-terminal amino acid sequences (van de Velde HJ et al (1994) J Cell Sci 107:2403-2416). Evidence suggests that NSP-A and NSP-C expression is restricted to neuronal and endocrine cell populations (van de Velde, supra). Immunohistochemical studies showed that rat NSP-A is expressed throughout the rat brain (van
15 de Velde HJ et al (1994) Mol Brain Res 23:81-92). NSP-B, however, is found only in a small cell lung carcinoma cell line and probably represents an aberrant NSP gene product (Roebroek AJ et al (1993) J Biol Chem 268:13439-13447). A previously reported neuronally expressed rat gene, CI-13, and two partially sequenced human cDNAs (GI 391043 and GI
20 894620), have a high degree of homology to NSPs which suggests that NSPs belong to a larger family of proteins (Wieczorek DF et al (1991) Mol Brain Res 10:33-41; Bell GI et al (1993) Hum Mol Genet 2:1793-798; Martin-Galla A et al (1992) Nat Genet 1:34-39).

25 Two large hydrophobic regions characterize the NSPs and homologous proteins and suggest membrane association. In fact, immunofluorescence and biochemical studies have established an association between NSPs and membranes of the ER (Senden NH et al (1994) Eur J Cell Biol 65:341-353). Analysis of NSP-A deletion mutants indicates that the carboxy-terminal hydrophobic region is necessary for membrane binding (van de Velde et al,
30 supra). Carboxy-terminal amino acid sequences of the NSPs are highly homologous, although they are not a perfect match to a consensus motif sufficient for retention of transmembrane proteins in the ER (van de Velde, supra; Jackson MR et al (1993) J Cell Biol 121:317-333). Thus, it appears likely that NSPs and related proteins are targeted to the ER by
35 conserved carboxy-terminal amino acids.

Immunostaining with anti-NSP-A antibodies suggests that NSP-A may be associated with both the rough and smooth neuronal ER. On the basis of this evidence and knowledge of neuronal ER function, van de Velde et al (1994; supra) conclude that NSPs may be involved in the protein
40 transport process or in the regulation of intracellular calcium levels in neuronal cells.

NSP-like Proteins and Disease

45 Dysfunction of ER-mediated neuronal protein transport may contribute to neurodegenerative diseases. For example, in amyotrophic lateral sclerosis (ALS), a degenerative disease of motor neurons,

position of neurofilaments in neuronal axons leads to dramatic defects in ER-mediated axonal transport of a variety of proteins (Collard JF et al (1995) Nature 375:61-64). Defects in protein transport have been further implicated in the pathogenesis of ALS by a transgenic mouse study in which ALS is modeled by a mutation in superoxide dismutase (SOD). SOD mutant animals displayed clinical and pathological features of human ALS and showed axonal transport defects associated with dilation of the ER (Mourelatos Z et al (1996) Proc Natl Acad Sci 93:5472-5477).

Analysis of specimens of a wide variety of primary human tumors show that NSP-A and NSP-C are expressed in small cell lung carcinoma, carcinoid tumors of the lung, but not in non-neuroendocrine non-small cell lung carcinomas (van de Velde et al (1994) Cancer Res 54:4769-4776). Furthermore, antibodies generated to small-cell lung carcinoma surface antigens recognize NSP-A, NSP-B, and NSP-C. Therefore, NSPs may act as markers in human lung cancer diagnosis and provide an avenue for corrective treatment (Senden NH et al (1994) Int J Cancer Suppl 8:84-88).

New NSP-like proteins could satisfy a need in the art by providing new means of diagnosing and treating cancer and neurodegenerative disorders such as ALS.

DISCLOSURE OF THE INVENTION

The present invention discloses two novel human NSP-like proteins (hereinafter referred to individually as NSPLPA and NSPLPB, and collectively as NSPLP), characterized as having homology to human NSP-A (GI 307307), NSP-B (GI 307309), NSP-C (GI 307311), and rat CI-13 (GI 281046). Accordingly, the invention features two substantially purified NSP-like proteins, as shown in amino acid sequence of SEQ ID NO:1 and SEQ ID NO:3, and having characteristics of NSPs.

One aspect of the invention features isolated and substantially purified polynucleotides which encode NSPLP. In a particular aspect, the polynucleotide is the nucleotide sequence of SEQ ID NO:2 or SEQ ID NO:4. In addition, the invention features polynucleotide sequences that hybridize under stringent conditions to SEQ ID NO:2 or SEQ ID NO:4.

The invention further relates to nucleic acid sequences encoding NSPLP, oligonucleotides, peptide nucleic acids (PNA), fragments, portions or antisense molecules thereof, and expression vectors and host cells comprising polynucleotides which encode NSPLP. The present invention also relates to antibodies which bind specifically to NSPLP, pharmaceutical compositions comprising substantially purified NSPLP, fragments thereof, or antagonists of NSPLP, in conjunction with a suitable pharmaceutical carrier, and methods for producing NSPLP, fragments thereof, or antagonists of NSPLP.

BRIEF DESCRIPTION OF DRAWINGS

Figures 1A, 1B and 1C show the amino acid sequence (SEQ ID NO:1) and nucleic acid sequence (SEQ ID NO:2) of the novel NSP-like protein,

NSPLPA. The alignment was produced using MacDNAsis software (Hitachi Software Engineering Co Ltd).

Figures 2A, 2B and 2C show the amino acid sequence (SEQ ID NO:3) and nucleic acid sequence (SEQ ID NO:4) of the novel NSP-like protein, NSPLPB (MacDNAsis software, Hitachi Software Engineering Co Ltd).

Figures 3A, 3B, 3C, 3D and 3E show the northern analysis for the consensus sequence (SEQ ID NO:4). The northern analysis was produced electronically using LIFESEQ™ database (Incyte Pharmaceuticals, Palo Alto CA).

Figures 4A, 4B and 4C show the northern analysis for Incyte Clones 31870 (SEQ ID NO:2) (LIFESEQ™ database, Incyte Pharmaceuticals, Palo Alto CA).

Figure 5 shows the assembly for the consensus sequence (SEQ ID NO:2).

Figures 6A, 6B, 6C, 6D, 6E and 6F show the amino acid sequence alignments among NSPLPA (SEQ ID NO:1), NSPLPB (SEQ ID NO:3), NSP-A (GI 307307; SEQ ID NO:5), NSP-B (GI 307309; SEQ ID NO:6), NSP-C (GI 307311; SEQ ID NO:7), and rat CI-13 (GI 281046 SEQ ID NO:8) produced using the multisequence alignment program of DNASTar software (DNASTar Inc, Madison WI).

Figure 7 shows the hydrophobicity plot (generated using MacDNAsis software) for NSPLPA, SEQ ID NO:1; the X axis reflects amino acid position, and the negative Y axis, hydrophobicity (Figs. 7, 8, and 9).

Figure 8 shows the hydrophobicity plot for NSPLPB, SEQ ID NO:3.

Figure 9 shows the hydrophobicity plot for NSP-C, SEQ ID NO:7.

MODES FOR CARRYING OUT THE INVENTION

Definitions

"Nucleic acid sequence" as used herein refers to an oligonucleotide, nucleotide or polynucleotide, and fragments or portions thereof, and to DNA or RNA of genomic or synthetic origin which may be single- or double-stranded, and represent the sense or antisense strand. Similarly, amino acid sequence as used herein refers to peptide or protein sequence.

"Peptide nucleic acid" as used herein refers to a molecule which comprises an oligomer to which an amino acid residue, such as lysine, and an amino group have been added. These small molecules, also designated anti-gene agents, stop transcript elongation by binding to their complementary (template) strand of nucleic acid (Nielsen PE et al (1993) Anticancer Drug Des 8:53-63).

As used herein, NSPLP refers to the amino acid sequences of substantially purified NSPLP obtained from any species, particularly mammalian, including bovine, ovine, porcine, murine, equine, and preferably human, from any source whether natural, synthetic, semi-synthetic or recombinant.

A "variant" of NSPLP is defined as an amino acid sequence that is altered by one or more amino acids. The variant may have "conservative" changes, wherein a substituted amino acid has similar structural or chemical properties, eg, replacement of leucine with isoleucine. More rarely, a variant may have "nonconservative" changes, eg, replacement of a glycine with a tryptophan. Similar minor variations may also include amino acid deletions or insertions, or both. Guidance in determining which and how many amino acid residues may be substituted, inserted or deleted without abolishing biological or immunological activity may be found using computer programs well known in the art, for example, DNASTar software.

A "deletion" is defined as a change in either amino acid or nucleotide sequence in which one or more amino acid or nucleotide residues, respectively, are absent.

An "insertion" or "addition" is that change in an amino acid or nucleotide sequence which has resulted in the addition of one or more amino acid or nucleotide residues, respectively, as compared to the naturally occurring NSPLP.

A "substitution" results from the replacement of one or more amino acids or nucleotides by different amino acids or nucleotides, respectively.

The term "biologically active" refers to a NSPLP having structural, regulatory or biochemical functions of a naturally occurring NSPLP. Likewise, "immunologically active" defines the capability of the natural, recombinant or synthetic NSPLP, or any oligopeptide thereof, to induce a specific immune response in appropriate animals or cells and to bind with specific antibodies.

The term "derivative" as used herein refers to the chemical modification of a nucleic acid encoding NSPLP or the encoded NSPLP. Illustrative of such modifications would be replacement of hydrogen by an alkyl, acyl, or amino group. A nucleic acid derivative would encode a polypeptide which retains essential biological characteristics of natural NSPLP.

As used herein, the term "substantially purified" refers to molecules, either nucleic or amino acid sequences, that are removed from their natural environment, isolated or separated, and are at least 60% free, preferably 75% free, and most preferably 90% free from other components with which they are naturally associated.

"Stringency" typically occurs in a range from about $T_m - 5^\circ\text{C}$ (5°C below the T_m of the probe) to about 20°C to 25°C below T_m . As will be understood by those of skill in the art, a stringency hybridization can be used to identify or detect identical polynucleotide sequences or to identify or detect similar or related polynucleotide sequences.

The term "hybridization" as used herein shall include "any process

by which a strand of nucleic acid joins with a complementary strand through base pairing" (Coombs J (1994) Dictionary of Biotechnology, Stockton Press, New York NY). Amplification as carried out in the polymerase chain reaction technologies is described in Dieffenbach CW and GS Dveksler (1995, PCR Primer, a Laboratory Manual, Cold Spring Harbor Press, Plainview NY).

Preferred Embodiments

The present invention relates to novel NSPLP and to the use of the nucleic acid and amino acid sequences in the study, diagnosis, prevention and treatment of disease. cDNAs encoding a portion of NSPLP were found in neuronal and endocrine tissue-derived cDNA libraries and in a variety of other tissues, including many types of tumors (Figs. 3A-3E and 4A-4C).

The present invention also encompasses NSPLP variants. A preferred NSPLP variant is one having at least 80% amino acid sequence similarity to the NSPLP amino acid sequence (SEQ ID NO:1), a more preferred NSPLP variant is one having at least 90% amino acid sequence similarity to SEQ ID NO:1 and a most preferred NSPLP variant is one having at least 95% amino acid sequence similarity to SEQ ID NO:1.

Nucleic acids encoding the human NSPLP of the present invention were first identified in cDNA, Incyte Clones 31870 (SEQ ID NO:4; THP-1 cell cDNA library, THP1NOB01) and 28742 (SEQ ID NO:9; fetal spleen cDNA library, SPLNFET01), through a computer-generated search for amino acid sequence alignments. A consensus sequence, SEQ ID NO:2, was derived from the following overlapping nucleic acid sequences: Incyte Clones 28742 (from cDNA library SPLNFET01); 45022, 45074, and 45509 (CORNNOT01); 121581 (MUSCNOT01); 570122 (MMLR3DT01); and 754150 (BRATUT02; Fig. 5). The nucleic acid sequence of SEQ ID NO:2 encodes the NSPLPA amino acid sequence, SEQ ID NO:1. The nucleic acid sequence of SEQ ID NO:4 encodes the NSPLPB amino acid sequence, SEQ ID NO:3. The nucleic acid sequence of SEQ ID NO:4 from residue C₄₉₆ to T₁₀₈ has 97% identity to the partial cDNA sequence of clone hbc043 (GI 39104; Bell et al, supra).

The present invention is based, in part, on the chemical and structural homology among NSPLPA, NSPLPB, NSP-A (GI 307307; Roebroek et al, supra), NSP-B (GI 307309; Roebroek et al, supra), NSP-C (GI 307311; Roebroek et al, supra), and rat CI-13 (GI 281046; Wieczorek et al, supra; Figs. 6A-D). NSPLPA and NSP-C share 66% identity, NSPLPB and NSP-C share 48% identity, while NSPLPA and NSPLPB share 50% identity. As illustrated by Figures 7, 8, and 9, NSPLPA, NSPLPB, and NSP-C have similar hydrophobicity plots suggesting similar structure. Like the NSPs, NSPLPA and NSPLPB have two large hydrophobic regions that could be used for membrane attachment. The carboxy-terminal amino acids Lys₁₉₉ through Lys₁₉₇ of NSPLPA precisely match, in position as well as sequence, an ER retention motif defined by Jackson et al (1993; supra). The novel NSPLPA is 199 amino acids long and has one potential N glycosylation site. The

novel NSPLPB is 241 amino acids long.

The NSPLP Coding Sequences

Th nucleic acid and deduced amino acid sequences of NSPLP are shown in Figures 1A, 1B, 1C, 2A, 2B and 2C. In accordance with the invention, any nucleic acid sequence which encodes the amino acid sequence of NSPLP can be used to generate recombinant molecules which express NSPLP. In a specific embodiment described herein, a nucleotide sequence encoding a portion of NSPLP was first isolated as Incyte Clones 31870 from a THP-1 cell cDNA library (THP1NOB01). While, Incyte Clone 28742 was first isolated from a fetal spleen cDNA library (SPLNFET01).

It will be appreciated by those skilled in the art that as a result of the degeneracy of the genetic code, a multitude of NSPLP-encoding nucleotide sequences, some bearing minimal homology to the nucleotide sequences of any known and naturally occurring gene may be produced. The invention contemplates each and every possible variation of nucleotide sequence that could be made by selecting combinations based on possible codon choices. These combinations are made in accordance with the standard triplet genetic code as applied to the nucleotide sequence of naturally occurring NSPLP, and all such variations are to be considered as being specifically disclosed.

Although nucleotide sequences which encode NSPLP and its variants are preferably capable of hybridizing to the nucleotide sequence of the naturally occurring NSPLP under appropriately selected conditions of stringency, it may be advantageous to produce nucleotide sequences encoding NSPLP or its derivatives possessing a substantially different codon usage. Codons may be selected to increase the rate at which expression of the peptide occurs in a particular prokaryotic or eukaryotic expression host in accordance with the frequency with which particular codons are utilized by the host. Other reasons for substantially altering the nucleotide sequence encoding NSPLP and its derivatives without altering the encoded amino acid sequences include the production of RNA transcripts having more desirable properties, such as a greater half-life, than transcripts produced from the naturally occurring sequence.

It is now possible to produce a DNA sequence, or portions thereof, encoding a NSPLP and its derivatives entirely by synthetic chemistry, after which the synthetic gene may be inserted into any of the many available DNA vectors and cell systems using reagents that are well known in the art at the time of the filing of this application. Moreover, synthetic chemistry may be used to introduce mutations into a sequence encoding NSPLP or any portion thereof.

Also included within the scope of the present invention are polynucleotide sequences that are capable of hybridizing to the nucleotide sequences of Figures 1A, 1B, 1C, 2A, 2B, and 2C under various

conditions of stringency. Hybridization conditions are based on the melting temperature (T_m) of the nucleic acid binding complex or probe, as taught in Berger and Kimmel (1987, Guide to Molecular Cloning Techniques, Methods in Enzymology, Vol 152, Academic Press, San Diego CA) incorporated herein by reference, and confer may be used at a defined stringency.

Altered nucleic acid sequences encoding NSPLP which may be used in accordance with the invention include deletions, insertions or substitutions of different nucleotides resulting in a polynucleotide that encodes the same or a functionally equivalent NSPLP. The protein may also show deletions, insertions or substitutions of amino acid residues which produce a silent change and result in a functionally equivalent NSPLP. Deliberate amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues as long as the biological activity of NSPLP is retained. For example, negatively charged amino acids include aspartic acid and glutamic acid; positively charged amino acids include lysine and arginine; and amino acids with uncharged polar head groups having similar hydrophilicity values include leucine, isoleucine, valine; glycine, alanine; asparagine, glutamine; serine, threonine phenylalanine, and tyrosine.

Included within the scope of the present invention are alleles of NSPLP. As used herein, an "allele" or "allelic sequence" is an alternative form of NSPLP. Alleles result from a mutation, ie, a change in the nucleic acid sequence, and generally produce altered mRNAs or polypeptides whose structure or function may or may not be altered. Any given gene may have none, one or many allelic forms. Common mutational changes which give rise to alleles are generally ascribed to natural deletions, additions or substitutions of amino acids. Each of these types of changes may occur alone, or in combination with the others, one or more times in a given sequence.

Methods for DNA sequencing are well known in the art and employ such enzymes as the Klenow fragment of DNA polymerase I, Sequenase® (US Biochemical Corp, Cleveland OH), Taq polymerase (Perkin Elmer, Norwalk CT), thermostable T7 polymerase (Amersham, Chicago IL), or combinations of recombinant polymerases and proofreading exonucleases such as the ELONGASE Amplification System marketed by Gibco BRL (Gaithersburg MD). Preferably, the process is automated with machines such as the Hamilton Micro Lab 2200 (Hamilton, Reno NV), Peltier Thermal Cycler (PTC200; MJ Research, Watertown MA) and the ABI 377 DNA sequencers (Perkin Elmer).

Extending the Polynucleotide Sequence

The polynucleotide sequence encoding NSPLP may be extended utilizing partial nucleotide sequence and various methods known in the art to detect upstream sequences such as promoters and regulatory

elements. Gobinda et al (1993; PCR Methods Applic 2:318-22) disclose "restriction-site" polymerase chain reaction (PCR) as a direct method which uses universal primers to retrieve unknown sequence adjacent to a known locus. First, genomic DNA is amplified in the presence of primer to a linker sequence and a primer specific to the known region. The amplified sequences are subjected to a second round of PCR with the same linker primer and another specific primer internal to the first one. Products of each round of PCR are transcribed with an appropriate RNA polymerase and sequenced using reverse transcriptase.

Inverse PCR can be used to amplify or extend sequences using divergent primers based on a known region (Triglia T et al (1988) Nucleic Acids Res 16:8186). The primers may be designed using OLIGO® 4.06 Primer Analysis Software (1992; National Biosciences Inc, Plymouth MN), or another appropriate program, to be 22-30 nucleotides in length, to have a GC content of 50% or more, and to anneal to the target sequence at temperatures about 68°-72° C. The method uses several restriction enzymes to generate a suitable fragment in the known region of a gene. The fragment is then circularized by intramolecular ligation and used as a PCR template.

Capture PCR (Lagerstrom M et al (1991) PCR Methods Applic 1:111-19) is a method for PCR amplification of DNA fragments adjacent to a known sequence in human and yeast artificial chromosome DNA. Capture PCR also requires multiple restriction enzyme digestions and ligations to place an engineered double-stranded sequence into an unknown portion of the DNA molecule before PCR.

Another method which may be used to retrieve unknown sequences is that of Parker JD et al (1991; Nucleic Acids Res 19:3055-60). Additionally, one can use PCR, nested primers and PromoterFinder libraries to walk in genomic DNA (PromoterFinder™ Clontech (Palo Alto CA). This process avoids the need to screen libraries and is useful in finding intron/exon junctions.

Preferred libraries for screening for full length cDNAs are ones that have been size-selected to include larger cDNAs. Also, random primed libraries are preferred in that they will contain more sequences which contain the 5' and upstream regions of genes. A randomly primed library may be particularly useful if an oligo d(T) library does not yield a full-length cDNA. Genomic libraries are useful for extension into the 5' nontranslated regulatory region.

Capillary electrophoresis may be used to analyze the size or confirm the nucleotide sequence of sequencing or PCR products. Systems for rapid sequencing are available from Perkin Elmer, Beckman Instruments (Fullerton CA), and other companies. Capillary sequencing may employ flowable polymers for electrophoretic separation, four different fluorescent dyes (one for each nucleotide) which are laser activated, and

detection of the emitted wavelengths by a charge coupled device camera. Output/light intensity is converted to electrical signal using appropriate software (eg. Genotyper™ and Sequence Navigator™ from Perkin Elmer) and the entire process from loading of samples to computer analysis and electronic data display is computer controlled. Capillary electrophoresis is particularly suited to the sequencing of small pieces of DNA which might be present in limited amounts in a particular sample. The reproducible sequencing of up to 350 bp of M13 phage DNA in 30 min has been reported (Ruiz-Martinez MC et al (1993) Anal Chem 65:2851-2858).

Expression of the Nucleotide Sequence

In accordance with the present invention, polynucleotide sequences which encode NSPLP, fragments of the polypeptide, fusion proteins or functional equivalents thereof may be used in recombinant DNA molecules that direct the expression of NSPLP in appropriate host cells. Due to the inherent degeneracy of the genetic code, other DNA sequences which encode substantially the same or a functionally equivalent amino acid sequence, may be used to clone and express NSPLP. As will be understood by those of skill in the art, it may be advantageous to produce NSPLP-encoding nucleotide sequences possessing non-naturally occurring codons. Codons preferred by a particular prokaryotic or eukaryotic host (Murray E et al (1989) Nuc Acids Res 17:477-508) can be selected, for example, to increase the rate of NSPLP expression or to produce recombinant RNA transcripts having desirable properties, such as a longer half-life, than transcripts produced from naturally occurring sequence.

The nucleotide sequences of the present invention can be engineered in order to alter a NSPLP coding sequence for a variety of reasons, including but not limited to, alterations which modify the cloning, processing and/or expression of the gene product. For example, mutations may be introduced using techniques which are well known in the art, eg, site-directed mutagenesis to insert new restriction sites, to alter glycosylation patterns, to change codon preference, to produce splice variants, etc.

In another embodiment of the invention, a natural, modified or recombinant polynucleotides encoding NSPLP may be ligated to a heterologous sequence to encode a fusion protein. For example, for screening of peptide libraries for inhibitors of NSPLP activity, it may be useful to encode a chimeric NSPLP protein that is recognized by a commercially available antibody. A fusion protein may also be engineered to contain a cleavage site located between a NSPLP sequence and the heterologous protein sequence, so that the NSPLP may be cleaved and purified away from the heterologous moiety.

In an alternate embodiment of the invention, the coding sequence of NSPLP may be synthesized, whole or in part, using chemical methods well known in the art (see Caruthers MH et al (1980) Nuc Acids Res Symp Ser

215-23, Horn T et al (1980) Nuc Acids Res Symp Ser 225-32, etc).
Alternatively, the protein itself could be produced using chemical
methods to synthesize a NSPLP amino acid sequence, whole or in part. For
example, peptide synthesis can be performed using various solid-phase
5 techniques (Roberge JY et al (1995) Science 269:202-204) and automated
synthesis may be achieved, for example, using the ABI 431A Peptide
Synthesizer (Perkin Elmer) in accordance with the instructions provided
by the manufacturer.

The newly synthesized peptide can be substantially by preparative
10 high performance liquid chromatography (eg, Creighton (1983) Proteins,
Structures and Molecular Principles, WH Freeman and Co, New York NY).
The composition of the synthetic peptides may be confirmed by amino acid
analysis or sequencing (eg, the Edman degradation procedure; Creighton,
supra). Additionally the amino acid sequence of NSPLP, or any part
15 thereof, may be altered during direct synthesis and/or combined using
chemical methods with sequences from other proteins, or any part thereof,
to produce a variant polypeptide.

Expression Systems

In order to express a biologically active NSPLP, the nucleotide
20 sequence encoding NSPLP or its functional equivalent, is inserted into an
appropriate expression vector, ie, a vector which contains the necessary
elements for the transcription and translation of the inserted coding
sequence.

Methods which are well known to those skilled in the art can be
25 used to construct expression vectors containing a NSPLP coding sequence
and appropriate transcriptional or translational controls. These methods
include in vitro recombinant DNA techniques, synthetic techniques and in
vivo recombination or genetic recombination. Such techniques are
described in Sambrook et al (1989) Molecular Cloning, A Laboratory
30 Manual, Cold Spring Harbor Press, Plainview NY and Ausubel FM et al
(1989) Current Protocols in Molecular Biology, John Wiley & Sons, New
York NY.

A variety of expression vector/host systems may be utilized to
contain and express a NSPLP coding sequence. These include but are not
35 limited to microorganisms such as bacteria transformed with recombinant
bacteriophage, plasmid or cosmid DNA expression vectors; yeast
transformed with yeast expression vectors; insect cell systems infected
with virus expression vectors (eg, baculovirus); plant cell systems
transfected with virus expression vectors (eg, cauliflower mosaic virus,
40 CaMV; tobacco mosaic virus, TMV) or transformed with bacterial expression
vectors (eg, Ti or pBR322 plasmid); or animal cell systems.

The "control elements" or "regulatory sequences" of these systems
vary in their strength and specificities and are those nontranslated
regions of the vector, enhancers, promoters, and 3' untranslated regions,

which interact with host cellular proteins to carry out transcription and translation. Depending on the vector system and host utilized, any number of suitable transcription and translation elements, including constitutive and inducible promoters, may be used. For example, when
5 cloning in bacterial systems, inducible promoters such as the hybrid lacZ promoter of the Bluescript® phagemid (Stratagene, LaJolla CA) or pSport1 (Gibco BRL) and ptrp-lac hybrids and the like may be used. The baculovirus polyhedrin promoter may be used in insect cells. Promoters or enhancers derived from the genomes of plant cells (eg, heat shock,
10 RUBISCO; and storage protein genes) or from plant viruses (eg, viral promoters or leader sequences) may be cloned into the vector. In mammalian cell systems, promoters from the mammalian genes or from mammalian viruses are most appropriate. If it is necessary to generate a cell line that contains multiple copies of NSPLP, vectors based on SV40
15 or EBV may be used with an appropriate selectable marker.

In bacterial systems, a number of expression vectors may be selected depending upon the use intended for NSPLP. For example, when large quantities of NSPLP are needed for the induction of antibodies, vectors which direct high level expression of fusion proteins that are
20 readily purified may be desirable. Such vectors include, but are not limited to, the multifunctional *E. coli* cloning and expression vectors such as Bluescript® (Stratagene), in which the NSPLP coding sequence may be ligated into the vector in frame with sequences for the amino-terminal Met and the subsequent 7 residues of β -galactosidase so that a hybrid
25 protein is produced; pIN vectors (Van Heeke & Schuster (1989) J Biol Chem 264:5503-5509); and the like. pGEX vectors (Promega, Madison WI) may also be used to express foreign polypeptides as fusion proteins with glutathione S-transferase (GST). In general, such fusion proteins are soluble and can easily be purified from lysed cells by adsorption to
30 glutathione-agarose beads followed by elution in the presence of free glutathione. Proteins made in such systems are designed to include heparin, thrombin or factor XA protease cleavage sites so that the cloned polypeptide of interest can be released from the GST moiety at will.

In the yeast, *Saccharomyces cerevisiae*, a number of vectors
35 containing constitutive or inducible promoters such as alpha factor, alcohol oxidase and PGH may be used. For reviews, see Ausubel et al (supra) and Grant et al (1987) Methods in Enzymology 153:516-544.

In cases where plant expression vectors are used, the expression of a sequence encoding NSPLP may be driven by any of a number of promoters.
40 For example, viral promoters such as the 35S and 19S promoters of CaMV (Brisson et al (1984) Nature 310:511-514) may be used alone or in combination with the omega leader sequence from TMV (Takamatsu et al (1987) EMBO J 6:307-311). Alternatively, plant promoters such as the small subunit of RUBISCO (Coruzzi et al (1984) EMBO J 3:1671-1680;

Broglie et al (1984) Science 224:838-843); or heat shock promoters (Winter J and Sinibaldi RM (1991) Results Probl Cell Differ 17:85-105) may be used. These constructs can be introduced into plant cells by direct DNA transformation or pathogen-mediated transfection. For reviews of such techniques, see Hobbs S or Murry LE in McGraw Hill Yearbook of Science and Technology (1992) McGraw Hill New York NY, pp 191-196 or Weissbach and Weissbach (1988) Methods for Plant Molecular Biology, Academic Press, New York NY, pp 421-463.

An alternative expression system which could be used to express NSPLP is an insect system. In one such system, Autographa californica nuclear polyhedrosis virus (AcNPV) is used as a vector to express foreign genes in Spodoptera frugiperda cells or in Trichoplusia larvae. The NSPLP coding sequence may be cloned into a nonessential region of the virus, such as the polyhedrin gene, and placed under control of the polyhedrin promoter. Successful insertion of NSPLP will render the polyhedrin gene inactive and produce recombinant virus lacking coat protein coat. The recombinant viruses are then used to infect S. frugiperda cells or Trichoplusia larvae in which NSPLP is expressed (Smith et al (1983) J Virol 46:584; Engelhard EK et al (1994) Proc Nat Acad Sci 91:3224-7).

In mammalian host cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, a NSPLP coding sequence may be ligated into an adenovirus transcription/translation complex consisting of the late promoter and tripartite leader sequence. Insertion in a nonessential E1 or E3 region of the viral genome will result in a viable virus capable of expressing NSPLP in infected host cells (Logan and Shenk (1984) Proc Natl Acad Sci 81:3655-59). In addition, transcription enhancers, such as the rous sarcoma virus (RSV) enhancer, may be used to increase expression in mammalian host cells.

Specific initiation signals may also be required for efficient translation of a NSPLP sequence. These signals include the ATG initiation codon and adjacent sequences. In cases where NSPLP, its initiation codon and upstream sequences are inserted into the appropriate expression vector, no additional translational control signals may be needed. However, in cases where only coding sequence, or a portion thereof, is inserted, exogenous transcriptional control signals including the ATG initiation codon must be provided. Furthermore, the initiation codon must be in the correct reading frame to ensure transcription of the entire insert. Exogenous transcriptional elements and initiation codons can be of various origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of enhancers appropriate to the cell system in use (Scharf D et al (1994) Results Probl Cell Differ 20:125-62; Bittner et al (1987) Methods in Enzymol 153:516-544).

In addition, a host cell strain may be chosen for its ability to modulate the expression of the inserted sequences or to process the expressed protein in the desired fashion. Such modifications of the polypeptide include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation and acylation.

Post-translational processing which cleaves a "prepro" form of the protein may also be important for correct insertion, folding and/or function. Different host cells such as CHO, HeLa, MDCK, 293, WI38, etc have specific cellular machinery and characteristic mechanisms for such post-translational activities and may be chosen to ensure the correct modification and processing of the introduced, foreign protein.

For long-term, high-yield production of recombinant proteins, stable expression is preferred. For example, cell lines which stably express NSPLP may be transformed using expression vectors which contain viral origins of replication or endogenous expression elements and a selectable marker gene. Following the introduction of the vector, cells may be allowed to grow for 1-2 days in an enriched media before they are switched to selective media. The purpose of the selectable marker is to confer resistance to selection, and its presence allows growth and recovery of cells which successfully express the introduced sequences. Resistant clumps of stably transformed cells can be proliferated using tissue culture techniques appropriate to the cell type.

Any number of selection systems may be used to recover transformed cell lines. These include, but are not limited to, the herpes simplex virus thymidine kinase (Wigler M et al (1977) Cell 11:223-32) and adenine phosphoribosyltransferase (Lowy I et al (1980) Cell 22:817-23) genes which can be employed in tk- or aprt- cells, respectively. Also, antimetabolite, antibiotic or herbicide resistance can be used as the basis for selection; for example, dhfr which confers resistance to methotrexate (Wigler M et al (1980) Proc Natl Acad Sci 77:3567-70); npt, which confers resistance to the aminoglycosides neomycin and G-418 (Colbere-Garapin F et al (1981) J Mol Biol 150:1-14) and als or pat, which confer resistance to chlorsulfuron and phosphinotricin acetyltransferase, respectively (Murry, supra). Additional selectable genes have been described, for example, trpB, which allows cells to utilize indole in place of tryptophan, or hisD, which allows cells to utilize histinol in place of histidine (Hartman SC and RC Mulligan (1988) Proc Natl Acad Sci 85:8047-51). Recently, the use of visible markers has gained popularity with such markers as anthocyanins, β glucuronidase and its substrate, GUS, and luciferase and its substrate, luciferin, being widely used not only to identify transformants, but also to quantify the amount of transient or stable protein expression attributable to a specific vector system (Rhodes CA et al (1995) Methods Mol Biol 55:121-131).

Identification of Transformants Containing the Polynucleotide Sequence

Although the presence/absence of marker gene expression suggests that the gene of interest is also present, its presence and expression should be confirmed. For example, if the NSPLP is inserted within a marker gene sequence, recombinant cells containing NSPLP can be identified by the absence of marker gene function. Alternatively, a marker gene can be placed in tandem with a NSPLP sequence under the control of a single promoter. Expression of the marker gene in response to induction or selection usually indicates expression of the tandem NSPLP as well.

Alternatively, host cells which contain the coding sequence for NSPLP and express NSPLP may be identified by a variety of procedures known to those of skill in the art. These procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridization and protein bioassay or immunoassay techniques which include membrane, solution, or chip based technologies for the detection and/or quantification of the nucleic acid or protein.

The presence of the polynucleotide sequence encoding NSPLP can be detected by DNA-DNA or DNA-RNA hybridization or amplification using probes, portions or fragments of polynucleotides encoding NSPLP. Nucleic acid amplification based assays involve the use of oligonucleotides or oligomers based on the NSPLP-encoding sequence to detect transformants containing DNA or RNA encoding NSPLP. As used herein "oligonucleotides" or "oligomers" refer to a nucleic acid sequence of at least about 10 nucleotides and as many as about 60 nucleotides, preferably about 15 to 30 nucleotides, and more preferably about 20-25 nucleotides which can be used as a probe or primer.

A variety of protocols for detecting and measuring the expression of NSPLP, using either polyclonal or monoclonal antibodies specific for the protein are known in the art. Examples include enzyme-linked immunosorbent assay (ELISA), radioimmunoassay (RIA) and fluorescent activated cell sorting (FACS). A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering epitopes on NSPLP is preferred, but a competitive binding assay may be employed. These and other assays are described, among other places, in Hampton R et al (1990, Serological Methods, a Laboratory Manual, APS Press, St Paul MN) and Maddox DE et al (1983, J Exp Med 158:1211).

A wide variety of labels and conjugation techniques are known by those skilled in the art and can be used in various nucleic acid and amino acid assays. Means for producing labeled hybridization or PCR probes for detecting sequences related to polynucleotides encoding NSPLP include oligolabeling, nick translation, end-labeling or PCR amplification using a labeled nucleotide. Alternatively, the NSPLP sequence, or any portion of it, may be cloned into a vector for the

production of an mRNA probe. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes in vitro by addition of an appropriate RNA polymerase such as T7, T3 or SP6 and labeled nucleotides.

5 A number of companies such as Pharmacia Biotech (Piscataway NJ), Promega (Madison WI), and US Biochemical Corp (Cleveland OH) supply commercial kits and protocols for these procedures. Suitable reporter molecules or labels include those radionuclides, enzymes, fluorescent, chemiluminescent, or chromogenic agents as well as substrates, cofactors,
10 inhibitors, magnetic particles and the like. Patents teaching the use of such labels include US Patents 3,817,837; 3,850,752; 3,939,350; 3,996,345; 4,277,437; 4,275,149 and 4,366,241. Also, recombinant immunoglobulins may be produced as shown in US Patent No. 4,816,567 incorporated herein by reference.

15 Purification of NSPLP

Host cells transformed with a nucleotide sequence encoding NSPLP may be cultured under conditions suitable for the expression and recovery of the encoded protein from cell culture. The protein produced by a recombinant cell may be secreted or contained intracellularly depending
20 on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing polynucleotides encoding NSPLP can be designed with signal sequences which direct secretion of NSPLP through a prokaryotic or eukaryotic cell membrane. Other recombinant constructions may join NSPLP to nucleotide sequence
25 encoding a polypeptide domain which will facilitate purification of soluble proteins (Kroll DJ et al (1993) DNA Cell Biol 12:441-53; cf discussion of vectors infra containing fusion proteins).

NSPLP may also be expressed as a recombinant protein with one or more additional polypeptide domains added to facilitate protein
30 purification. Such purification facilitating domains include, but are not limited to, metal chelating peptides such as histidine-tryptophan modules that allow purification on immobilized metals, protein A domains that allow purification on immobilized immunoglobulin, and the domain utilized in the FLAGS extension/affinity purification system (Immunex
35 Corp, Seattle WA). The inclusion of a cleavable linker sequences such as Factor XA or enterokinase (Invitrogen, San Diego CA) between the purification domain and NSPLP is useful to facilitate purification. One such expression vector provides for expression of a fusion protein
40 comprising an NSPLP and contains nucleic acid encoding 6 histidine residues followed by thioredoxin and an enterokinase cleavage site. The histidine residues facilitate purification on IMIAC (immobilized metal ion affinity chromatography as described in Porath et al (1992) Protein Expression and Purification 3: 263-281) while the enterokinase cleavage site provides a means for purifying NSPLP from the fusion protein.

In addition to recombinant production, fragments of NSPLP may be produced by direct peptide synthesis using solid-phase techniques (cf Stewart et al (1969) Solid-Phase Peptide Synthesis, WH Freeman Co, San Francisco; Merrifield J (1963) J Am Chem Soc 85:2149-2154). In vitro protein synthesis may be performed using manual techniques or by automation. Automated synthesis may be achieved, for example, using Applied Biosystems 431A Peptide Synthesizer (Perkin Elmer, Foster City CA) in accordance with the instructions provided by the manufacturer. Various fragments of NSPLP may be chemically synthesized separately and combined using chemical methods to produce the full length molecule.

Uses of NSPLP

The rationale for use of the nucleotide and polypeptide sequences disclosed herein is based in part on the chemical and structural homology among the novel NSPLP proteins disclosed herein, NSP-A (GI 307307; Roebroek et al, supra), NSP-B (GI 307309; Roebroek et al, supra), NSP-C (GI 307311; Roebroek et al, supra), and rat CI-13 (GI 281046; Wieczorek et al, supra).

Accordingly, NSPLP or a NSPLP derivative may be used to treat cancer and neurodegenerative disorders, such as ALS. In those conditions where NSPLP protein activity is not desirable, cells could be transfected with antisense sequences of NSPLP-encoding polynucleotides or provided with antagonists of NSPLP.

NSPLP Antibodies

NSPLP-specific antibodies are useful for the diagnosis of conditions and diseases associated with expression of NSPLP. Such antibodies may include, but are not limited to, polyclonal, monoclonal, chimeric, single chain, Fab fragments and fragments produced by a Fab expression library. Neutralizing antibodies, ie, those which inhibit dimer formation, are especially preferred for diagnostics and therapeutics.

NSPLP for antibody induction does not require biological activity; however, the protein fragment, or oligopeptide must be antigenic. Peptides used to induce specific antibodies may have an amino acid sequence consisting of at least five amino acids, preferably at least 10 amino acids. Preferably, they should mimic a portion of the amino acid sequence of the natural protein and may contain the entire amino acid sequence of a small, naturally occurring molecule. Short stretches of NSPLP amino acids may be fused with those of another protein such as keyhole limpet hemocyanin and antibody produced against the chimeric molecule. Procedures well known in the art can be used for the production of antibodies to NSPLP.

For the production of antibodies, various hosts including goats, rabbits, rats, mice, etc may be immunized by injection with NSPLP or any portion, fragment or oligopeptide which retains immunogenic properties.

Depending on the host species, various adjuvants may be used to increase immunological response. Such adjuvants include but are not limited to, Freund's, mineral gels such as aluminum hydroxide, and surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, keyhole limpet hemocyanin, and dinitrophenol. BCG (bacilli Calmette-Guerin) and Corynebacterium parvum are potentially useful human adjuvants.

Monoclonal antibodies to NSPLP may be prepared using any technique which provides for the production of antibody molecules by continuous cell lines in culture. These include but are not limited to the hybridoma technique originally described by Koehler and Milstein (1975 Nature 256:495-497), the human B-cell hybridoma technique (Kosbor et al (1983) Immunol Today 4:72; Cote et al (1983) Proc Natl Acad Sci 80:2026-2030) and the EBV-hybridoma technique (Cole et al (1985) Monoclonal Antibodies and Cancer Therapy, Alan R Liss Inc, New York NY, pp 77-96).

In addition, techniques developed for the production of "chimeric antibodies", the splicing of mouse antibody genes to human antibody genes to obtain a molecule with appropriate antigen specificity and biological activity can be used (Morrison et al (1984) Proc Natl Acad Sci 81:6851-6855; Neuberger et al (1984) Nature 312:604-608; Takeda et al (1985) Nature 314:452-454). Alternatively, techniques described for the production of single chain antibodies (US Patent No. 4,946,778) can be adapted to produce NSPLP-specific single chain antibodies

Antibodies may also be produced by inducing in vivo production in the lymphocyte population or by screening recombinant immunoglobulin libraries or panels of highly specific binding reagents as disclosed in Orlandi et al (1989, Proc Natl Acad Sci 86: 3833-3837), and Winter G and Milstein C (1991; Nature 349:293-299).

Antibody fragments which contain specific binding sites for NSPLP may also be generated. For example, such fragments include, but are not limited to, the F(ab')₂ fragments which can be produced by pepsin digestion of the antibody molecule and the Fab fragments which can be generated by reducing the disulfide bridges of the F(ab')₂ fragments. Alternatively, Fab expression libraries may be constructed to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity (Huse WD et al (1989) Science 256:1275-1281).

A variety of protocols for competitive binding or immunoradiometric assays using either polyclonal or monoclonal antibodies with established specificities are well known in the art. Such immunoassays typically involve the formation of complexes between NSPLP and its specific antibody and the measurement of complex formation. A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two noninterfering epitopes on a specific NSPLP protein is preferred, but

a competitive binding assay may also be employed. These assays are described in Maddox DE et al (1983, J Exp Med 158:1211).

Diagnostic Assays Using NSPLP Specific Antibodies

Particular NSPLP antibodies are useful for the diagnosis of conditions or diseases characterized by expression of NSPLP or in assays to monitor patients being treated with NSPLP, agonists or inhibitors. Diagnostic assays for NSPLP include methods utilizing the antibody and a label to detect NSPLP in human body fluids or extracts of cells or tissues. The polypeptides and antibodies of the present invention may be used with or without modification. Frequently, the polypeptides and antibodies will be labeled by joining them, either covalently or noncovalently, with a reporter molecule. A wide variety of reporter molecules are known, several of which were described above.

A variety of protocols for measuring NSPLP, using either polyclonal or monoclonal antibodies specific for the respective protein are known in the art. Examples include enzyme-linked immunosorbent assay (ELISA), radioimmunoassay (RIA) and fluorescent activated cell sorting (FACS). A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering epitopes on NSPLP is preferred, but a competitive binding assay may be employed. These assays are described, among other places, in Maddox, DE et al (1983, J Exp Med 158:1211).

In order to provide a basis for diagnosis, normal or standard values for NSPLP expression must be established. This is accomplished by combining body fluids or cell extracts taken from normal subjects, either animal or human, with antibody to NSPLP under conditions suitable for complex formation which are well known in the art. The amount of standard complex formation may be quantified by comparing various artificial membranes containing known quantities of NSPLP with both control and disease samples from biopsied tissues. Then, standard values obtained from normal samples may be compared with values obtained from samples from subjects potentially affected by disease. Deviation between standard and subject values establishes the presence of disease state.

Drug Screening

NSPLP, its catalytic or immunogenic fragments or oligopeptides thereof, can be used for screening therapeutic compounds in any of a variety of drug screening techniques. The fragment employed in such a test may be free in solution, affixed to a solid support, borne on a cell surface, or located intracellularly. The formation of binding complexes, between NSPLP and the agent being tested, may be measured.

Another technique for drug screening which may be used provides for high throughput screening of compounds having suitable binding affinity to the NSPLP is described in detail in "Determination of Amino Acid Sequence Antigenicity" by Geysen HN, WO Application 84/03564, published on September 13, 1984, and incorporated herein by reference. In summary,

large numbers of different small peptide test compounds are synthesized on a solid substrate, such as plastic pins or some other surface. The peptide test compounds are reacted with fragments of NSPLP and washed. Bound NSPLP is then detected by methods well known in the art. Purified NSPLP can also be coated directly onto plates for use in the aforementioned drug screening techniques. Alternatively, non-neutralizing antibodies can be used to capture the peptide and immobilize it on a solid support.

This invention also contemplates the use of competitive drug screening assays in which neutralizing antibodies capable of binding NSPLP specifically compete with a test compound for binding NSPLP. In this manner, the antibodies can be used to detect the presence of any peptide which shares one or more antigenic determinants with NSPLP.

Uses of the Polynucleotide Encoding NSPLP

A polynucleotide encoding NSPLP, or any part thereof, may be used for diagnostic and/or therapeutic purposes. For diagnostic purposes, polynucleotides encoding NSPLP of this invention may be used to detect and quantitate gene expression in biopsied tissues in which expression of NSPLP may be implicated. The diagnostic assay is useful to distinguish between absence, presence, and excess expression of NSPLP and to monitor regulation of NSPLP levels during therapeutic intervention. Included in the scope of the invention are oligonucleotide sequences, antisense RNA and DNA molecules, and PNAs.

Another aspect of the subject invention is to provide for hybridization or PCR probes which are capable of detecting polynucleotide sequences, including genomic sequences, encoding NSPLP or closely related molecules. The specificity of the probe, whether it is made from a highly specific region, eg, 10 unique nucleotides in the 5' regulatory region, or a less specific region, eg, especially in the 3' region, and the stringency of the hybridization or amplification (maximal, high, intermediate or low) will determine whether the probe identifies only naturally occurring sequences encoding NSPLP, alleles or related sequences.

Probes may also be used for the detection of related sequences and should preferably contain at least 50% of the nucleotides from any of these NSPLP encoding sequences. The hybridization probes of the subject invention may be derived from the nucleotide sequence of SEQ ID NO:2 or from genomic sequence including promoter, enhancer elements and introns of the naturally occurring NSPLP. Hybridization probes may be labeled by a variety of reporter groups, including radionuclides such as ³²P or ³⁵S, or enzymatic labels such as alkaline phosphatase coupled to the probe via avidin/biotin coupling systems, and the like.

Other means for producing specific hybridization probes for DNAs encoding NSPLP include the cloning of nucleic acid sequences encoding

NSPLP or NSPLP derivatives into vectors for the production of mRNA probes. Such vectors are known in the art and are commercially available and may be used to synthesize RNA probes in vitro by means of the addition of the appropriate RNA polymerase as T7 or SP6 RNA polymerase and the appropriate radioactively labeled nucleotides.

Polynucleotide sequences encoding NSPLP may be used for the diagnosis of conditions or diseases with which the expression of NSPLP is associated. For example, polynucleotide sequences encoding NSPLP may be used in hybridization or PCR assays of fluids or tissues from biopsies to detect NSPLP expression. The form of such qualitative or quantitative methods may include Southern or northern analysis, dot blot or other membrane-based technologies; PCR technologies; dip stick, pIN, chip and ELISA technologies. All of these techniques are well known in the art and are the basis of many commercially available diagnostic kits.

The nucleotide sequences encoding NSPLP disclosed herein provide the basis for assays that detect activation or induction associated with cancer and neurodegenerative disorders, such as ALS. The nucleotide sequence encoding NSPLP may be labeled by methods known in the art and added to a fluid or tissue sample from a patient under conditions suitable for the formation of hybridization complexes. After an incubation period, the sample is washed with a compatible fluid which optionally contains a dye (or other label requiring a developer) if the nucleotide has been labeled with an enzyme. After the compatible fluid is rinsed off, the dye is quantitated and compared with a standard. If the amount of dye in the biopsied or extracted sample is significantly elevated over that of a comparable control sample, the nucleotide sequence has hybridized with nucleotide sequences in the sample, and the presence of elevated levels of nucleotide sequences encoding NSPLP in the sample indicates the presence of the associated disease.

Such assays may also be used to evaluate the efficacy of a particular therapeutic treatment regime in animal studies, in clinical trials, or in monitoring the treatment of an individual patient. In order to provide a basis for the diagnosis of disease, a normal or standard profile for NSPLP expression must be established. This is accomplished by combining body fluids or cell extracts taken from normal subjects, either animal or human, with NSPLP, or a portion thereof, under conditions suitable for hybridization or amplification. Standard hybridization may be quantified by comparing the values obtained for normal subjects with a dilution series of NSPLP run in the same experiment where a known amount of a substantially purified NSPLP is used. Standard values obtained from normal samples may be compared with values obtained from samples from patients afflicted with NSPLP-associated diseases. Deviation between standard and subject values is used to establish the presence of disease.

Once disease is established, a therapeutic agent is administered and a treatment profile is generated. Such assays may be repeated on a regular basis to evaluate whether the values in the profile progress toward or return to the normal or standard pattern. Successive treatment profiles may be used to show the efficacy of treatment over a period of several days or several months.

PCR, as described in US Patent Nos. 4,683,195 and 4,965,188, provides additional uses for oligonucleotides based upon the NSPLP sequence. Such oligomers are generally chemically synthesized, but they may be generated enzymatically or produced from a recombinant source. Oligomers generally comprise two nucleotide sequences, one with sense orientation (5'→3') and one with antisense (3'←5'), employed under optimized conditions for identification of a specific gene or condition. The same two oligomers, nested sets of oligomers, or even a degenerate pool of oligomers may be employed under less stringent conditions for detection and/or quantitation of closely related DNA or RNA sequences.

Additionally, methods which may be used to quantitate the expression of a particular molecule include radiolabeling (Melby PC et al 1993 J Immunol Methods 159:235-44) or biotinylating (Duplaa C et al 1993 Anal Biochem 229-36) nucleotides, coamplification of a control nucleic acid, and standard curves onto which the experimental results are interpolated. Quantitation of multiple samples may be speeded up by running the assay in an ELISA format where the oligomer of interest is presented in various dilutions and a spectrophotometric or colorimetric response gives rapid quantitation. For example, the presence of a relatively high amount of NSPLP in extracts of biopsied tissues may indicate the onset of cancer. A definitive diagnosis of this type may allow health professionals to begin aggressive treatment and prevent further worsening of the condition. Similarly, further assays can be used to monitor the progress of a patient during treatment. Furthermore, the nucleotide sequences disclosed herein may be used in molecular biology techniques that have not yet been developed, provided the new techniques rely on properties of nucleotide sequences that are currently known such as the triplet genetic code, specific base pair interactions, and the like.

Therapeutic Use

Based upon its homology to genes encoding NSP-like proteins and its expression profile, polynucleotide sequences encoding NSPLP disclosed herein may be useful in the treatment of conditions such as cancer and neurodegenerative disorders, such as ALS.

Expression vectors derived from retroviruses, adenovirus, herpes or vaccinia viruses, or from various bacterial plasmids, may be used for delivery of nucleotide sequences to the targeted organ, tissue or cell population. Methods which are well known to those skilled in the art can

be used to construct recombinant vectors which will express antisense polynucleotides of the gene encoding NSPLP. See, for example, the techniques described in Sambrook et al (supra) and Ausubel et al (supra).

The polynucleotides comprising full length cDNA sequence and/or its regulatory elements enable researchers to use sequences encoding NSPLP as an investigative tool in sense (Yousoufian H and HF Lodish 1993 Mol Cell Biol 13:98-104) or antisense (Eguchi et al (1991) Annu Rev Biochem 60:631-652) regulation of gene function. Such technology is now well known in the art, and sense or antisense oligomers, or larger fragments, can be designed from various locations along the coding or control regions.

Genes encoding NSPLP can be turned off by transfecting a cell or tissue with expression vectors which express high levels of a desired NSPLP-encoding fragment. Such constructs can flood cells with untranslatable sense or antisense sequences. Even in the absence of integration into the DNA, such vectors may continue to transcribe RNA molecules until all copies are disabled by endogenous nucleases. Transient expression may last for a month or more with a non-replicating vector (Mettler I, personal communication) and even longer if appropriate replication elements are part of the vector system.

As mentioned above, modifications of gene expression can be obtained by designing antisense molecules, DNA, RNA or PNA, to the control regions of gene encoding NSPLP, ie, the promoters, enhancers, and introns. Oligonucleotides derived from the transcription initiation site, eg, between -10 and +10 regions of the leader sequence, are preferred. The antisense molecules may also be designed to block translation of mRNA by preventing the transcript from binding to ribosomes. Similarly, inhibition can be achieved using "triple helix" base-pairing methodology. Triple helix pairing compromises the ability of the double helix to open sufficiently for the binding of polymerases, transcription factors, or regulatory molecules. Recent therapeutic advances using triplex DNA were reviewed by Gee JE et al (In: Huber BE and BI Carr (1994) Molecular and Immunologic Approaches, Futura Publishing Co, Mt Kisco NY).

Ribozymes are enzymatic RNA molecules capable of catalyzing the specific cleavage of RNA. The mechanism of ribozyme action involves sequence-specific hybridization of the ribozyme molecule to complementary target RNA, followed by endonucleolytic cleavage. Within the scope of the invention are engineered hammerhead motif ribozyme molecules that can specifically and efficiently catalyze endonucleolytic cleavage of sequences encoding NSPLP.

Specific ribozyme cleavage sites within any potential RNA target are initially identified by scanning the target molecule for ribozyme cleavage sites which include the following sequences, GUA, GUU and GUC.

Once identified, short RNA sequences of between 15 and 20 ribonucleotides corresponding to the region of the target gene containing the cleavage site may be evaluated for secondary structural features which may render the oligonucleotide inoperable. The suitability of candidate targets may also be evaluated by testing accessibility to hybridization with complementary oligonucleotides using ribonuclease protection assays.

Antisense molecules and ribozymes of the invention may be prepared by any method known in the art for the synthesis of RNA molecules. These include techniques for chemically synthesizing oligonucleotides such as solid phase phosphoramidite chemical synthesis. Alternatively, RNA molecules may be generated by *in vitro* and *in vivo* transcription of DNA sequences encoding NSPLP. Such DNA sequences may be incorporated into a wide variety of vectors with suitable RNA polymerase promoters such as T7 or SP6. Alternatively, antisense cDNA constructs that synthesize antisense RNA constitutively or inducibly can be introduced into cell lines, cells or tissues.

RNA molecules may be modified to increase intracellular stability and half-life. Possible modifications include, but are not limited to, the addition of flanking sequences at the 5' and/or 3' ends of the molecule or the use of phosphorothioate or 2' O-methyl rather than phosphodiesterase linkages within the backbone of the molecule. This concept is inherent in the production of PNAs and can be extended in all of these molecules by the inclusion of nontraditional bases such as inosine, queosine and wybutosine as well as acetyl-, methyl-, thio- and similarly modified forms of adenine, cytidine, guanine, thymine, and uridine which are not as easily recognized by endogenous endonucleases.

Methods for introducing vectors into cells or tissues include those methods discussed *infra* and which are equally suitable for *in vivo*, *in vitro* and *ex vivo* therapy. For *ex vivo* therapy, vectors are introduced into stem cells taken from the patient and clonally propagated for autologous transplant back into that same patient is presented in US Patent Nos. 5,399,493 and 5,437,994, disclosed herein by reference. Delivery by transfection and by liposome are quite well known in the art.

Furthermore, the nucleotide sequences for NSPLP disclosed herein may be used in molecular biology techniques that have not yet been developed, provided the new techniques rely on properties of nucleotide sequences that are currently known, including but not limited to such properties as the triplet genetic code and specific base pair interactions.

Detection and Mapping of Related Polynucleotide Sequences

The nucleic acid sequence for NSPLP can also be used to generate hybridization probes for mapping the naturally occurring genomic sequence. The sequence may be mapped to a particular chromosome or to a specific region of the chromosome using well known techniques. These

include in situ hybridization to chromosomal spreads, flow-sorted chromosomal preparations, or artificial chromosome constructions such as yeast artificial chromosomes, bacterial artificial chromosomes, bacterial P1 constructions or single chromosome cDNA libraries as reviewed in Price CM (1993; Blood Rev 7:127-34) and Trask BJ (1991; Trends Genet 7:149-54).

The technique of fluorescent in situ hybridization of chromosome spreads has been described, among other places, in Verma et al (1988) Human Chromosomes: A Manual of Basic Techniques, Pergamon Press, New York NY. Fluorescent in situ hybridization of chromosomal preparations and other physical chromosome mapping techniques may be correlated with additional genetic map data. Examples of genetic map data can be found in the 1994 Genome Issue of Science (265:1981f). Correlation between the location of the gene encoding NSPLP on a physical chromosomal map and a specific disease (or predisposition to a specific disease) may help delimit the region of DNA associated with that genetic disease. The nucleotide sequences of the subject invention may be used to detect differences in gene sequences between normal, carrier or affected individuals.

In situ hybridization of chromosomal preparations and physical mapping techniques such as linkage analysis using established chromosomal markers may be used for extending genetic maps. For example an sequence tagged site based map of the human genome was recently published by the Whitehead-MIT Center for Genomic Research (Hudson TJ et al(1995) Science 270:1945-1954). Often the placement of a gene on the chromosome of another mammalian species such as mouse (Whitehead Institute/MIT Center for Genome Research, Genetic Map of the Mouse, Database Release 10, April 28, 1995) may reveal associated markers even if the number or arm of a particular human chromosome is not known. New sequences can be assigned to chromosomal arms, or parts thereof, by physical mapping. This provides valuable information to investigators searching for disease genes using positional cloning or other gene discovery techniques. Once a disease or syndrome, such as ataxia telangiectasia (AT), has been crudely localized by genetic linkage to a particular genomic region, for example, AT to 11q22-23 (Gatti et al (1988) Nature 336:577-580), any sequences mapping to that area may represent associated or regulatory genes for further investigation. The nucleotide sequence of the subject invention may also be used to detect differences in the chromosomal location due to translocation, inversion, etc. among normal, carrier or affected individuals.

Pharmaceutical Compositions

The present invention relates to pharmaceutical compositions which may comprise nucleotides, proteins, antibodies, agonists, antagonists, or inhibitors, alone or in combination with at least one other agent, such as stabilizing compound, which may be administered in any sterile,

biocompatible pharmaceutical carrier, including, but not limited to, saline, buffered saline, dextrose, and water. Any of these molecules can be administered to a patient alone, or in combination with other agents, drugs or hormones, in pharmaceutical compositions where it is mixed with excipient(s) or pharmaceutically acceptable carriers. In one embodiment of the present invention, the pharmaceutically acceptable carrier is pharmaceutically inert.

Administration of Pharmaceutical Compositions

Administration of pharmaceutical compositions is accomplished orally or parenterally. Methods of parenteral delivery include topical, intra-arterial (directly to the tumor), intramuscular, subcutaneous, intramedullary, intrathecal, intraventricular, intravenous, intraperitoneal, or intranasal administration. In addition to the active ingredients, these pharmaceutical compositions may contain suitable pharmaceutically acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically. Further details on techniques for formulation and administration may be found in the latest edition of "Remington's Pharmaceutical Sciences" (Maack Publishing Co, Easton PA).

Pharmaceutical compositions for oral administration can be formulated using pharmaceutically acceptable carriers well known in the art in dosages suitable for oral administration. Such carriers enable the pharmaceutical compositions to be formulated as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions and the like, for ingestion by the patient.

Pharmaceutical preparations for oral use can be obtained through combination of active compounds with solid excipient, optionally grinding a resulting mixture, and processing the mixture of granules, after adding suitable auxiliaries, if desired, to obtain tablets or dragee cores. Suitable excipients are carbohydrate or protein fillers such as sugars, including lactose, sucrose, mannitol, or sorbitol; starch from corn, wheat, rice, potato, or other plants; cellulose such as methyl cellulose, hydroxypropylmethyl-cellulose, or sodium carboxymethylcellulose; and gums including arabic and tragacanth; and proteins such as gelatin and collagen. If desired, disintegrating or solubilizing agents may be added, such as the cross-linked polyvinyl pyrrolidone, agar, alginic acid, or a salt thereof, such as sodium alginate.

Dragee cores are provided with suitable coatings such as concentrated sugar solutions, which may also contain gum arabic, talc, polyvinylpyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may be added to the tablets or dragee coatings for product identification or to characterize the quantity of

active compound, ie, dosage.

Pharmaceutical preparations which can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a coating such as glycerol or sorbitol. Push-fit capsules
5 can contain active ingredients mixed with a filler or binders such as lactose or starches, lubricants such as talc or magnesium stearate, and, optionally, stabilizers. In soft capsules, the active compounds may be dissolved or suspended in suitable liquids, such as fatty oils, liquid paraffin, or liquid polyethylene glycol with or without stabilizers.

10 Pharmaceutical formulations for parenteral administration include aqueous solutions of active compounds. For injection, the pharmaceutical compositions of the invention may be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hanks's solution, Ringer's solution, or physiologically buffered saline. Aqueous
15 injection suspensions may contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Additionally, suspensions of the active compounds may be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils such as sesame oil, or synthetic
20 fatty acid esters, such as ethyl oleate or triglycerides, or liposomes. Optionally, the suspension may also contain suitable stabilizers or agents which increase the solubility of the compounds to allow for the preparation of highly concentrated solutions.

For topical or nasal administration, penetrants appropriate to the
25 particular barrier to be permeated are used in the formulation. Such penetrants are generally known in the art.

Manufacture and Storage

The pharmaceutical compositions of the present invention may be
30 manufactured in a manner that known in the art, eg, by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping or lyophilizing processes.

The pharmaceutical composition may be provided as a salt and can be
35 formed with many acids, including but not limited to hydrochloric, sulfuric, acetic, lactic, tartaric, malic, succinic, etc. Salts tend to be more soluble in aqueous or other protonic solvents that are the corresponding free base forms. In other cases, the preferred preparation may be a lyophilized powder in 1mM-50 mM histidine, 0.1%-2% sucrose,
2%-7% mannitol at a pH range of 4.5 to 5.5 that is combined with buffer prior to use.

40 After pharmaceutical compositions comprising a compound of the invention formulated in a acceptable carrier have been prepared, they can be placed in an appropriate container and labeled for treatment of an indicated condition. For administration of NSPLP, such labeling would include amount, frequency and method of administration.

Therapeutically Effective Dose

Pharmaceutical compositions suitable for use in the present invention include compositions wherein the active ingredients are contained in an effective amount to achieve the intended purpose. The determination of an effective dose is well within the capability of those skilled in the art.

For any compound, the therapeutically effective dose can be estimated initially either in cell culture assays, eg, of neoplastic cells, or in animal models, usually mice, rabbits, dogs, or pigs. The animal model is also used to achieve a desirable concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans.

A therapeutically effective dose refers to that amount of protein or its antibodies, antagonists, or inhibitors which ameliorate the symptoms or condition. Therapeutic efficacy and toxicity of such compounds can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, eg, ED50 (the dose therapeutically effective in 50% of the population) and LD50 (the dose lethal to 50% of the population). The dose ratio between therapeutic and toxic effects is the therapeutic index, and it can be expressed as the ratio, LD50/ED50. Pharmaceutical compositions which exhibit large therapeutic indices are preferred. The data obtained from cell culture assays and animal studies is used in formulating a range of dosage for human use. The dosage of such compounds lies preferably within a range of circulating concentrations that include the ED50 with little or no toxicity. The dosage varies within this range depending upon the dosage form employed, sensitivity of the patient, and the route of administration.

The exact dosage is chosen by the individual physician in view of the patient to be treated. Dosage and administration are adjusted to provide sufficient levels of the active moiety or to maintain the desired effect. Additional factors which may be taken into account include the severity of the disease state, eg, tumor size and location; age, weight and gender of the patient; diet, time and frequency of administration, drug combination(s), reaction sensitivities, and tolerance/response to therapy. Long acting pharmaceutical compositions might be administered every 3 to 4 days, every week, or once every two weeks depending on half-life and clearance rate of the particular formulation.

Normal dosage amounts may vary from 0.1 to 100,000 micrograms, up to a total dose of about 1 g, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature and generally available to practitioners in the art. Those skilled in the art will employ different formulations for nucleotides than for proteins or their inhibitors. Similarly, delivery of polynucleotides or polypeptides will be specific to particular cells,

conditions, locations, etc.

It is contemplated, for example, that NSPLP or an NSPLP derivative can be delivered in a suitable formulation to block the progression of cancerous cell growth or of neuronal degeneration. Similarly, administration of NSPLP antagonists may also inhibit the activity or shorten the lifespan of this protein.

The examples below are provided to illustrate the subject invention and are not included for the purpose of limiting the invention.

INDUSTRIAL APPLICABILITY

I Construction of cDNA Libraries

THP-1

THP-1 is a human leukemic cell line derived from the blood of a 1-year-old boy with acute monocytic leukemia. The THP-1 cells represent monocytes. The THP-1 cDNA library was custom constructed by Stratagene (Stratagene, 11099 M. Torrey Pines Rd., La Jolla, CA 92037) essentially as described below.

Stratagene prepared the cDNA library using oligo d(T) priming. Synthetic adapter oligonucleotides were ligated onto the cDNA molecules enabling them to be inserted into the Uni-ZAP™ vector system (Stratagene). This allowed high efficiency unidirectional (sense orientation) lambda library construction and the convenience of a plasmid system with blue/white color selection to detect clones with cDNA insertions.

The quality of the cDNA library was screened using DNA probes, and then, the pBluescript® phagemid (Stratagene) was excised. This phagemid allows the use of a plasmid system for easy insert characterization, sequencing, site-directed mutagenesis, the creation of unidirectional deletions and expression of fusion polypeptides. Subsequently, the custom-constructed library phage particles were infected into *E. coli* host strain XL1-Blue® (Stratagene). The high transformation efficiency of this bacterial strain increases the probability that the cDNA library will contain rare, under-represented clones. Alternative unidirectional vectors include, but are not limited to, pcDNAI (Invitrogen, San Diego CA) and pSHlox-1 (Novagen, Madison WI).

Fetal spleen

The human spleen cell cDNA library was custom constructed by Stratagene (catalogue # 937205. Stratagene, La Jolla CA). The starting cell population is mixed, having been obtained from fetal spleens which have a diverse cell population. Furthermore, the fetal spleens have been pooled from different sources. Poly(A+) RNA (mRNA) was purified from the spleen cells. cDNA was synthesized from the mRNA. Synthetic adaptor oligonucleotides were ligated onto cDNA ends enabling its insertion into Uni-ZAP™ vector system (Stratagene), allowing high efficiency unidirectional (sense orientation) lambda library construction and the

convenience of a plasmid system with blue/white color selection to detect clones with cDNA insertions. Alternative unidirectional vectors are pcDNA1 (invitrogen, San Diego CA) and pSHlox-1 (Novagen, Madison WI).

II Isolation of cDNA Clones

THP-1

The phagemid forms of individual cDNA clones were obtained by the in vivo excision process, in which the host bacterial strain was co-infected with both the library phage and an f1 helper phage. Polypeptides or enzymes derived from both the library-containing phage and the helper phage nicked the DNA, initiated new DNA synthesis from defined sequences on the target DNA, and created a smaller, single stranded circular phagemid DNA molecule that included all DNA sequences of the pBluescript phagemid and the cDNA insert. The phagemid DNA was released from the cells and purified, and used to reinfect fresh host cells (SOLR, Stratagene) where double-stranded phagemid DNA was produced. Because the phagemid carries the gene for β -lactamase, the newly transformed bacteria were selected on medium containing ampicillin.

An alternate method of purifying phagemid has recently become available. It utilizes the Miniprep Kit (Catalog No. 77468, available from Advanced Genetic Technologies Corp., 19212 Orbit Drive, Gaithersburg, Maryland). This kit is in the 96-well format and provides enough reagents for 960 purifications. Each kit is provided with a recommended protocol, which has been employed except for the following changes. First, the 96 wells are each filled with only 1 ml of sterile terrific broth with carbenicillin at 25 mg/L and glycerol at 0.4%. After the wells are inoculated, the bacteria are cultured for 24 hours and lysed with 60 μ l of lysis buffer. A centrifugation step (2900 rpm for 5 minutes) is performed before the contents of the block are added to the primary filter plate. The optional step of adding isopropanol to TRIS buffer is not routinely performed. After the last step in the protocol, samples are transferred to a Beckman 96-well block for storage.

Phagemid DNA was also purified using the QIAWELL-8 Plasmid Purification System from the QIAGEN® DNA Purification System (QIAGEN Inc, Chatsworth CA). This product provides a convenient, rapid and reliable high-throughput method for lysing the bacterial cells and isolating highly purified phagemid DNA using QIAGEN anion-exchange resin particles with EMPORE™ membrane technology from 3M in a multiwell format. The DNA was eluted from the purification resin and prepared for DNA sequencing and other analytical manipulations.

Fetal spleen

The phagemid forms of individual cDNA clones were obtained by the in vivo excision process, in which the host bacterial strain was co-infected with both the library phage and an f1 helper phage. Polypeptides or enzymes derived from both the library-containing phage and the helper phage nicked the DNA, initiated new DNA synthesis from

defined sequences on the target DNA, and created a smaller, single stranded circular phagemid DNA molecule that included all DNA sequences of the pBluescript phagemid and the cDNA insert. The phagemid DNA was released from the cells and purified, and used to reinfect fresh host cells (SOLR, Stratagene) where double-stranded phagemid DNA was produced. Because the phagemid carries the gene for β -lactamase, the newly transformed bacteria were selected on medium containing ampicillin.

Phagemid DNA was also purified using the QIAWELL-8 Plasmid Purification System from the QIAGEN® DNA Purification System (QIAGEN Inc, Chatsworth CA). This product provides a convenient, rapid and reliable high-throughput method for lysing the bacterial cells and isolating highly purified phagemid DNA using QIAGEN anion-exchange resin particles with EMPORE™ membrane technology from 3M in a multiwell format. The DNA was eluted from the purification resin and prepared for DNA sequencing and other analytical manipulations.

III Homology Searching of cDNA Clones and Their Deduced Proteins

Each cDNA was compared to sequences in GenBank using a search algorithm developed by Applied Biosystems and incorporated into the INHERIT™ 670 Sequence Analysis System. In this algorithm, Pattern Specification Language (TRW Inc, Los Angeles CA) was used to determine regions of homology. The three parameters that determine how the sequence comparisons run were window size, window offset, and error tolerance. Using a combination of these three parameters, the DNA database was searched for sequences containing regions of homology to the query sequence, and the appropriate sequences were scored with an initial value. Subsequently, these homologous regions were examined using dot matrix homology plots to distinguish regions of homology from chance matches. Smith-Waterman alignments were used to display the results of the homology search.

Peptide and protein sequence homologies were ascertained using the INHERIT- 670 Sequence Analysis System in a way similar to that used in DNA sequence homologies. Pattern Specification Language and parameter windows were used to search protein databases for sequences containing regions of homology which were scored with an initial value. Dot-matrix homology plots were examined to distinguish regions of significant homology from chance matches.

BLAST, which stands for Basic Local Alignment Search Tool (Altschul SF (1993) J Mol Evol 36:290-300; Altschul, SF et al (1990) J Mol Biol 215:403-10), was used to search for local sequence alignments. BLAST produces alignments of both nucleotide and amino acid sequences to determine sequence similarity. Because of the local nature of the alignments, BLAST is especially useful in determining exact matches or in identifying homologs. BLAST is useful for matches which do not contain gaps. The fundamental unit of BLAST algorithm output is the High-scoring

Segment Pair (HSP).

An HSP consists of two sequence fragments of arbitrary but equal lengths whose alignment is locally maximal and for which the alignment score meets or exceeds a threshold or cutoff score set by the user. The BLAST approach is to look for HSPs between a query sequence and a database sequence, to evaluate the statistical significance of any matches found, and to report only those matches which satisfy the user-selected threshold of significance. The parameter E establishes the statistically significant threshold for reporting database sequence matches. E is interpreted as the upper bound of the expected frequency of chance occurrence of an HSP (or set of HSPs) within the context of the entire database search. Any database sequence whose match satisfies E is reported in the program output.

IV Northern Analysis

Northern analysis is a laboratory technique used to detect the presence of a transcript of a gene and involves the hybridization of a labelled nucleotide sequence to a membrane on which RNAs from a particular cell type or tissue have been bound (Sambrook et al. supra).

Analogous computer techniques using BLAST (Altschul SF 1993 and 1990, supra) are used to search for identical or related molecules in nucleotide databases such as GenBank or the LIFESEQ™ database (Incyte, Palo Alto CA). This analysis is much faster than multiple, membrane-based hybridizations. In addition, the sensitivity of the computer search can be modified to determine whether any particular match is categorized as exact or homologous.

The basis of the search is the product score which is defined as:

$$\frac{\% \text{ sequence identity} \times \% \text{ maximum BLAST score}}{100}$$

100

and it takes into account both the degree of similarity between two sequences and the length of the sequence match. For example, with a product score of 40, the match will be exact within a 1-2% error; and at 70, the match will be exact. Homologous molecules are usually identified by selecting those which show product scores between 15 and 40, although lower scores may identify related molecules.

V Extension of NSPLP-Encoding Polynucleotides to Full Length or to Recover Regulatory Elements

Full length NSPLP-encoding nucleic acid sequence (SEQ ID NO:2) is used to design oligonucleotide primers for extending a partial nucleotide sequence to full length or for obtaining 5' sequences from genomic libraries. One primer is synthesized to initiate extension in the antisense direction (XLR) and the other is synthesized to extend sequence in the sense direction (XLF). Primers allow the extension of the known NSPLP-encoding sequence "outward" generating amplicons containing new, unknown nucleotide sequence for the region of interest (US Patent

Application 08/487,112, filed June 7, 1995, specifically incorporated by reference). The initial primers are designed from the cDNA using OLIGO® 4.06 Primer Analysis Software (National Biosciences), or another appropriate program, to be 22-30 nucleotides in length, to have a GC content of 50% or more, and to anneal to the target sequence at temperatures about 68°-72° C. Any stretch of nucleotides which would result in hairpin structures and primer-primer dimerizations is avoided.

The original, selected cDNA libraries, or a human genomic library are used to extend the sequence; the latter is most useful to obtain 5' upstream regions. If more extension is necessary or desired, additional sets of primers are designed to further extend the known region.

By following the instructions for the XL-PCR kit (Perkin Elmer) and thoroughly mixing the enzyme and reaction mix, high fidelity amplification is obtained. Beginning with 40 pmol of each primer and the recommended concentrations of all other components of the kit, PCR is performed using the Peltier Thermal Cycler (PTC200; MJ Research, Watertown MA) and the following parameters:

Step 1	94° C for 1 min (initial denaturation)
Step 2	65° C for 1 min
Step 3	68° C for 6 min
Step 4	94° C for 15 sec
Step 5	65° C for 1 min
Step 6	68° C for 7 min
Step 7	Repeat step 4-6 for 15 additional cycles
Step 8	94° C for 15 sec
Step 9	65° C for 1 min
Step 10	68° C for 7:15 min
Step 11	Repeat step 8-10 for 12 cycles
Step 12	72° C for 8 min
Step 13	4° C (and holding)

A 5-10 µl aliquot of the reaction mixture is analyzed by electrophoresis on a low concentration (about 0.6-0.8%) agarose mini-gel to determine which reactions were successful in extending the sequence. Bands thought to contain the largest products were selected and cut out of the gel. Further purification involves using a commercial gel extraction method such as QIAQuick™ (QIAGEN Inc). After recovery of the DNA, Klenow enzyme was used to trim single-stranded, nucleotide overhangs creating blunt ends which facilitate religation and cloning.

After ethanol precipitation, the products are redissolved in 13 µl of ligation buffer, 1µl T4-DNA ligase (15 units) and 1µl T4 polynucleotide kinase are added, and the mixture is incubated at room temperature for 2-3 hours or overnight at 16° C. Competent *E. coli* cells (in 40 µl of appropriate media) are transformed with 3 µl of ligation mixture and cultured in 80 µl of SOC medium (Sambrook J et al, supra). After incubation for one hour at 37° C, the whole transformation mixture is plated on Luria Bertani (LB)-agar (Sambrook J et al, supra) containing 2xCarb. The following day, several colonies are randomly picked from each plate and cultured in 150 µl of liquid LB/2xCarb medium placed in an

individual well of an appropriate, commercially-available, sterile 96-well microtiter plate. The following day, 5 μ l of each overnight culture is transferred into a non-sterile 96-well plate and after dilution 1:10 with water, 5 μ l of each sample is transferred into a PCR array.

For PCR amplification, 18 μ l of concentrated PCR reaction mix (3.3x) containing 4 units of rTth DNA polymerase, a vector primer and one or both of the gene specific primers used for the extension reaction are added to each well. Amplification is performed using the following conditions:

Step 1	94° C for 60 sec
Step 2	94° C for 20 sec
Step 3	55° C for 30 sec
Step 4	72° C for 90 sec
Step 5	Repeat steps 2-4 for an additional 29 cycles
Step 6	72° C for 180 sec
Step 7	4° C (and holding)

Aliquots of the PCR reactions are run on agarose gels together with molecular weight markers. The sizes of the PCR products are compared to the original partial cDNAs, and appropriate clones are selected, ligated into plasmid and sequenced.

VI Labeling and Use of Hybridization Probes

Hybridization probes derived from SEQ ID NO:2 are employed to screen cDNAs, genomic DNAs or mRNAs. Although the labeling of oligonucleotides, consisting of about 20 base-pairs, is specifically described, essentially the same procedure is used with larger cDNA fragments. Oligonucleotides are designed using state-of-the-art software such as OLIGO 4.06 (National Biosciences), labeled by combining 50 pmol of each oligomer and 250 mCi of [γ -³²P] adenosine triphosphate (Amersham, Chicago IL) and T4 polynucleotide kinase (DuPont NEN[®], Boston MA). The labeled oligonucleotides are substantially purified with Sephadex G-25 super fine resin column (Pharmacia). A portion containing 10⁷ counts per minute of each of the sense and antisense oligonucleotides is used in a typical membrane based hybridization analysis of human genomic DNA digested with one of the following endonucleases (Ase I, Bgl II, Eco RI, Pst I, Xba I, or Pvu II; DuPont NEN[®]).

The DNA from each digest is fractionated on a 0.7 percent agarose gel and transferred to nylon membranes (Nytran Plus, Schleicher & Schuell, Durham NH). Hybridization is carried out for 16 hours at 40°C. To remove nonspecific signals, blots are sequentially washed at room temperature under increasingly stringent conditions up to 0.1 x saline sodium citrate and 0.5% sodium dodecyl sulfate. After XOMAT AR[™] film (Kodak, Rochester NY) is exposed to the blots in a Phosphorimager cassette (Molecular Dynamics, Sunnyvale CA) for several hours, hybridization patterns are compared visually.

VII Antisense Molecules

The NSPLP-encoding sequence, or any part thereof, is used to inhibit in vivo or in vitro expression of naturally occurring NSPLP. Although use of antisense oligonucleotides, comprising about 20 base-pairs, is specifically described, essentially the same procedure is used with larger cDNA fragments. An oligonucleotide based on the coding sequences of NSPLP, as shown in Figs. 1A, 1B, 2A, and 2B is used to inhibit expression of naturally occurring NSPLP. The complementary oligonucleotide is designed from the most unique 5' sequence as shown in Figures 1A, 1B, 2A, and 2B and used either to inhibit transcription by preventing promoter binding to the upstream nontranslated sequence or translation of an NSPLP-encoding transcript by preventing the ribosome from binding. Using an appropriate portion of the leader and 5' sequence of SEQ ID NO:2, an effective antisense oligonucleotide includes any 15-20 nucleotides spanning the region which translates into the signal or early coding sequence of the polypeptide as shown in Figures 1A, 1B, 2A, and 2B.

VIII Expression of NSPLP

Expression of the NSPLP is accomplished by subcloning the cDNAs into appropriate vectors and transfecting the vectors into host cells. In this case, the cloning vector, pSport, previously used for the generation of the cDNA library is used to express NSPLP in E. coli. Upstream of the cloning site, this vector contains a promoter for β -galactosidase, followed by sequence containing the amino-terminal Met and the subsequent 7 residues of β -galactosidase. Immediately following these eight residues is a bacteriophage promoter useful for transcription and a linker containing a number of unique restriction sites.

Induction of an isolated, transfected bacterial strain with IPTG using standard methods produces a fusion protein which consists of the first seven residues of β -galactosidase, about 5 to 15 residues of linker, and the full length NSPLP-encoding sequence. The signal sequence directs the secretion of NSPLP into the bacterial growth media which can be used directly in the following assay for activity.

IX NSPLP Activity

NSPLP's ER targeting activity can be assessed by a method of van de Velde et al (1994, supra). Microsomes are collected from cells expressing NSPLP by a 100,000 g spin in a method described by Verboomen H et al (1992 Biochem J 286:591-596). After treatment with 0.5 M KCl and centrifugation the pellet is resuspended and subject to gel electrophoresis. Western blot analysis using antibodies to NSPLP reveals the presence of NSPLP in the ER membrane.

X Production of NSPLP Specific Antibodies

NSPLP substantially purified using PAGE electrophoresis (Sambrook, supra) is used to immunize rabbits and to produce antibodies using standard protocols. The amino acid sequence translated from NSPLP is analyzed using DNASTar software (DNASTar Inc) to determine regions of high immunogenicity and a corresponding oligopolypeptide is synthesized and used to raise antibodies by means known to those of skill in the art. Analysis to select appropriate epitopes, such as those near the C-terminus or in hydrophilic regions (shown in Figs. 7 and 8) is described by Ausubel FM et al (supra).

Typically, the oligopeptides are 15 residues in length, synthesized using an Applied Biosystems Peptide Synthesizer Model 431A using fmoc-chemistry, and coupled to keyhole limpet hemocyanin (KLH, Sigma) by reaction with M-maleimidobenzoyl-N-hydroxysuccinimide ester (MBS; Ausubel FM et al, supra). Rabbits are immunized with the oligopeptide-KLH complex in complete Freund's adjuvant. The resulting antisera are tested for antipeptide activity, for example, by binding the peptide to plastic, blocking with 1% BSA, reacting with rabbit antisera, washing, and reacting with radioiodinated, goat anti-rabbit IgG.

XI Purification of Naturally Occurring NSPLP Using Specific Antibodies

Naturally occurring or recombinant NSPLP is substantially purified by immunoaffinity chromatography using antibodies specific for NSPLP. An immunoaffinity column is constructed by covalently coupling NSPLP antibody to an activated chromatographic resin such as CnBr-activated Sepharose (Pharmacia Biotech). After the coupling, the resin is blocked and washed according to the manufacturer's instructions.

Media containing NSPLP is passed over the immunoaffinity column, and the column is washed under conditions that allow the preferential absorbance of NSPLP (eg, high ionic strength buffers in the presence of detergent). The column is eluted under conditions that disrupt antibody/NSPLP binding (eg, a buffer of pH 2-3 or a high concentration of a chaotrope such as urea or thiocyanate ion), and NSPLP is collected.

XII Identification of Molecules Which Interact with NSPLP

NSPLP, or biologically active fragments thereof, are labelled with ¹²⁵I Bolton-Hunter reagent (Bolton, AE and Hunter, WM (1973) Biochem J 133: 529). Candidate molecules previously arrayed in the wells of a 96 well plate are incubated with the labelled NSPLP, washed and any wells with labelled NSPLP complex are assayed. Data obtained using different concentrations of NSPLP are used to calculate values for the number, affinity, and association of NSPLP with the candidate molecules.

All publications and patents mentioned in the above specification are herein incorporated by reference. Various modifications and variations of the described method and system of the invention will be apparent to those skilled in the art without departing from the scope and

spirit of the invention. Although the invention has been described in connection with specific preferred embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific embodiments. Indeed, various modifications of the described modes for carrying out the invention which are obvious to those skilled in molecular biology or related fields are intended to be within the scope of the following claims.

SEQUENCE LISTING

(1) GENERAL INFORMATION

- (i) APPLICANT: INCYTE PHARMACEUTICALS, INC.
- (ii) TITLE OF THE INVENTION: TWO NOVEL HUMAN NSP-LIKE PROTEINS
- (iii) NUMBER OF SEQUENCES: 9
- (iv) CORRESPONDENCE ADDRESS:
 - (A) ADDRESSEE: Incyte Pharmaceuticals, Inc.
 - (B) STREET: 3174 Porter Drive
 - (C) CITY: Palo Alto
 - (D) STATE: CA
 - (E) COUNTRY: U.S.
 - (F) ZIP: 94304
- (v) COMPUTER READABLE FORM:
 - (A) MEDIUM TYPE: Diskette
 - (B) COMPUTER: IBM Compatible
 - (C) OPERATING SYSTEM: DOS
 - (D) SOFTWARE: FastSEQ Version 1.5
- (vi) CURRENT APPLICATION DATA:
 - (A) PCT APPLICATION NUMBER: To Be Assigned
 - (B) FILING DATE: Filed Herewith
- (vii) PRIOR APPLICATION DATA:
 - (A) APPLICATION NUMBER: 08/700,607
 - (B) FILING DATE: AUGUST 12, 1996
- (viii) ATTORNEY/AGENT INFORMATION:
 - (A) NAME: Billings, Lucy J.
 - (B) REGISTRATION NUMBER: 36,749
 - (C) REFERENCE/DOCKET NUMBER: PF-0114 PCT
- (ix) TELECOMMUNICATION INFORMATION:
 - (A) TELEPHONE: 415-855-0555
 - (B) TELEFAX: 415-845-4166

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 199 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(vii) IMMEDIATE SOURCE:

- (A) LIBRARY:
- (B) CLONE: Consensus

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

```

Met Asp Gly Gln Lys Lys Asn Trp Lys Asp Lys Val Val Asp Leu Leu
 1           5           10           15
Tyr Trp Arg Asp Ile Lys Lys Thr Gly Val Val Phe Gly Ala Ser Leu
          20          25          30
Phe Leu Leu Leu Ser Leu Thr Val Phe Ser Ile Val Ser Val Thr Ala
 35          40          45
Tyr Ile Ala Leu Ala Leu Leu Ser Val Thr Ile Ser Phe Arg Ile Tyr
 50          55          60
Lys Gly Val Ile Gln Ala Ile Gln Lys Ser Asp Glu Gly His Pro Phe
 65          70          75          80
Arg Ala Tyr Leu Glu Ser Glu Val Ala Ile Ser Glu Glu Leu Val Gln
          85          90          95
Lys Tyr Ser Asn Ser Ala Leu Gly His Val Asn Cys Thr Ile Lys Glu
          100         105         110
Leu Arg Arg Leu Phe Leu Val Asp Leu Val Asp Ser Leu Lys Phe
          115         120         125
Ala Val Leu Met Trp Val Phe Thr Tyr Val Gly Ala Leu Phe Asn Gly
          130         135         140
Leu Thr Leu Leu Ile Leu Ala Leu Ile Ser Leu Phe Ser Val Pro Val
          145         150         155         160
Ile Tyr Glu Arg His Gln Ala Gln Ile Asp His Tyr Leu Gly Leu Ala
          165         170         175
Asn Lys Asn Val Lys Asp Ala Met Ala Lys Ile Gln Ala Lys Ile Pro
          180         185         190
Gly Leu Lys Arg Lys Ala Glu
          195

```

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 799 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vii) IMMEDIATE SOURCE:

- (A) LIBRARY:
- (B) CLONE: Consensus

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

```

GGTTTGTGCA GTTACAGCTT TTCTNTTGGT ATGCATAATT AATANTTGGG GCTGCAAAGA      60
GATCGTGACA AGAGATGGAC GGTCAGAAGA AAAATTGGAA GGACAAGGTT GTTGACCTCC      120
TGTA CTGGAG AGACATTAAG AAGACTGGAG TGGTGT TGG TGCCAGCCTA TTCCTGCTGC      180

```

TTTCATTGAC	AGTATTCAGC	ATTGTGAGCG	TAACAGCCTA	CATTGCCTTG	GCCCTGCTCT	240
CTGTGACCAT	CAGCTTTAGG	ATATACAAGG	GTGTGATCCA	AGCTATCCAG	AAATCAGATG	300
AAGGCCACCC	ATTCAGGGCA	TATCTGGAAT	CTGAAGTTGC	TATATCTGAG	GAGTTGGTTC	360
AGAAGTACAG	TAATTCTGCT	CTTGGTCATG	TGAACTGCAC	GATAAAGGAA	CTCAGGCGCC	420
TCTTCTTAGT	TGATGATTTA	GTTGATTCTC	TGAAGTTTGC	AGTGTTGATG	TGGGTATTTA	480
CCTATGTTGG	TGCCTTGTTT	AATGGTCTGA	CACTACTGAT	TTTGGCTCTC	ATTTCACTCT	540
TCAGTGTTCC	TGTTATTTAT	GAACGGCATC	AGGCACAGAT	AGATCATTAT	CTAGGACTTG	600
CAAATAAGAA	TGTTAAAGAT	GCTATGGCTA	AAATCCAAGC	AAAAATCCCT	GGATTGAAGC	660
GCAAAGCTGA	ATGAAAACGC	CCAAAATAAT	TAGTAGGAGT	TCATCTTTAA	AGGGGATATT	720
CATTTGATTA	TACGGGGGAG	GGTCAGGGAA	GAACGACCTT	GACGTTGCAG	TGCAGTTTCA	780
CAGATCGTTG	TTAGATCTT					799

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 241 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(vii) IMMEDIATE SOURCE:

- (A) LIBRARY: THP1NOB01
- (B) CLONE: 31870

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

Met	Ala	Glu	Arg	Xaa	Ala	Ala	Thr	Gln	Ser	His	Ser	Ile	Ser	Ser	Ser	1	5	10	15
Ser	Phe	Gly	Ala	Glu	Pro	Ser	Ala	Pro	Gly	Gly	Gly	Gly	Ser	Pro	Gly	20	25	30	
Ala	Cys	Pro	Ala	Leu	Gly	Thr	Lys	Ser	Cys	Ser	Ser	Ser	Cys	Ala	Val	35	40	45	
His	Asp	Leu	Ile	Xaa	Trp	Arg	Asp	Val	Lys	Lys	Thr	Gly	Phe	Val	Phe	50	55	60	
Gly	Thr	Thr	Leu	Ile	Met	Leu	Leu	Ser	Leu	Ala	Ala	Phe	Ser	Val	Ile	65	70	75	80
Ser	Val	Val	Ser	Tyr	Leu	Ile	Leu	Ala	Leu	Leu	Ser	Val	Thr	Ile	Ser	85	90	95	
Phe	Arg	Ile	Tyr	Lys	Ser	Val	Ile	Gln	Ala	Val	Gln	Lys	Ser	Glu	Glu	100	105	110	
Gly	His	Pro	Phe	Lys	Ala	Tyr	Leu	Asp	Val	Asp	Ile	Thr	Leu	Ser	Ser	115	120	125	
Glu	Ala	Phe	His	Asn	Tyr	Met	Asn	Ala	Ala	Met	Val	His	Ile	Asn	Arg	130	135	140	
Ala	Leu	Lys	Leu	Ile	Ile	Arg	Leu	Phe	Leu	Val	Glu	Asp	Leu	Val	Asp	145	150	155	160
Ser	Leu	Lys	Leu	Ala	Val	Phe	Met	Trp	Leu	Met	Thr	Tyr	Val	Gly	Ala	165	170	175	
Val	Phe	Asn	Gly	Ile	Thr	Leu	Leu	Ile	Leu	Ala	Glu	Leu	Leu	Ile	Xaa	180	185	190	
Ser	Val	Pro	Ile	Val	Tyr	Xaa	Lys	Tyr	Lys	Val	Pro	Ser	Lys	Thr	Pro	195	200	205	
Trp	Asn	Arg	Gln	Lys	Lys	Gly	Arg	Ile	Ser	Thr	Trp	Lys	Pro	Glu	Met	210	215	220	
Gln	Gln	Leu	Leu	Lys	His	His	Leu	Ile	Val	Ile	Thr	Ser	Leu	Leu	Val	225	230	235	240
Leu																			

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1095 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vii) IMMEDIATE SOURCE:

- (A) LIBRARY: THP1NOB01
 (B) CLONE: 31870

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

ACACNAGCGN	NTCGNGCTCC	CGAACCTCTA	GCTGCGACTC	GGANTGAGTC	AGTCAGTCTG	60
TCGGAGTCTG	TCCTCGGAGC	AGGCGGAGTA	AAGGGACTTG	AGCGAGCCAG	TTGCCGGATT	120
ATTCTATTTT	CCCTCCCTCT	CTCCCGCCCC	GTATCTCTTT	TCATTTTNNT	NCCACCCTTG	180
CTCGCGTANC	ATGGCGGAGC	GTNCGGCGGC	CACTCAGTCC	CATTCCATCT	CCTCGTCGTC	240
CTTCGGAGCC	GAGCCGTCCG	CGCCCGGCGG	CGGCGGGAGC	CCAGGAGCCT	GCCCCGCCCT	300
GGGGACGAAG	AGCTGCAGCT	CCTCCTGTGC	GGTGCACGAT	CTGATTTTMT	GGAGAGATGT	360
GAAGAAGACT	GGGTTTGTCT	TTGGCACCAC	GCTGATCATG	CTGCTTTCCC	TGGCAGCTTT	420
CAGTGTGATC	AGTGTGGTTT	CTTACCTCAT	CCTGGCTCTT	CTCTCTGTCA	CCATCAGCTT	480
CAGGATCTAC	AAGTCCGTCA	TCCAAGCTGT	ACAGAAGTCA	GAAGAAGGCC	ATCCATTCAA	540
AGCCTACCTG	GACGTAGACA	TTACTCTGTC	CTCAGAAGCT	TTCCATAATT	ACATGAATGC	600
TGCCATGGTG	CACATCAACA	GGGCCCTGAA	ACTCATTATT	CGTCTCTTTC	TGGTAGAAGA	660
TCTGGTTGAC	TCCTTGAAGC	TGGCTGTCTT	CATGTGGCTG	ATGACCTATG	TTGGTGCTGT	720
TTTTAACGGA	ATCACCTTTC	TAATTCTTGC	TGAACTGCTC	ATTTTNAGTG	TCCCGATTGT	780
NTATNAGAAG	TACAAGGTTT	CAAGCAAAAC	TCCCTGGAAT	CGCCAAAAAA	AAGGCAGAAT	840
AAGTACATGG	AAACCAGAAA	TGCAACAGTT	ACTAAAACAC	CATTTAATAG	TTATAACGTC	900
GTTACTTGTA	CTATGAAGGA	AAATACTCAG	TGTCAGCTTG	AGCCTGCATT	CCAAGCTTTT	960
TTTTTAATTT	GGTGGTTTTT	TCCCATCCTT	TCCCTTTAAC	CCTCAGTNTC	AAGCACAAAN	1020
TTTNTATGGAC	TGATAANNGA	TCTATNTTAG	ANCTCAGAAG	ANGANAGNTT	CANNTGCATA	1080
GGNTAAGGNA	NTACC					1095

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 776 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(vii) IMMEDIATE SOURCE:

- (A) LIBRARY: GenBank
 (B) CLONE: 307307

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

Met	Ala	Ala	Pro	Gly	Asp	Pro	Gln	Asp	Glu	Leu	Leu	Pro	Leu	Ala	Gly
1				5				10						15	
Pro	Gly	Ser	Gln	Trp	Leu	Arg	His	Arg	Gly	Glu	Gly	Glu	Asn	Glu	Ala
			20					25					30		
Val	Thr	Pro	Lys	Gly	Ala	Thr	Pro	Ala	Pro	Gln	Ala	Gly	Glu	Pro	Ser
			35				40					45			
Pro	Gly	Leu	Gly	Ala	Arg	Ala	Arg	Glu	Ala	Ala	Ser	Arg	Glu	Ala	Gly
			50			55					60				
Ser	Gly	Pro	Ala	Arg	Gln	Ser	Pro	Val	Ala	Met	Glu	Thr	Ala	Ser	Thr
					70					75				80	
Gly	Val	Ala	Gly	Val	Ser	Ser	Ala	Met	Asp	His	Thr	Phe	Ser	Thr	Thr
				85					90					95	

Ser	Lys	Asp	Gly	Glu	Gly	Ser	Cys	Tyr	Thr	Ser	Leu	Ile	Ser	Asp	Ile
			100					105					110		
Cys	Tyr	Pro	Pro	Gln	Glu	Asp	Ser	Thr	Tyr	Phe	Thr	Gly	Ile	Leu	Gln
		115					120					125			
Lys	Glu	Asn	Gly	His	Val	Thr	Ile	Ser	Glu	Ser	Pro	Glu	Glu	Leu	Gly
	130					135					140				
Thr	Pro	Gly	Pro	Ser	Leu	Pro	Asp	Val	Pro	Gly	Ile	Glu	Ser	Arg	Gly
145					150					155					160
Leu	Phe	Ser	Ser	Asp	Ser	Gly	Ile	Glu	Met	Thr	Pro	Ala	Glu	Ser	Thr
				165					170					175	
Glu	Val	Asn	Lys	Ile	Leu	Ala	Asp	Pro	Leu	Asp	Gln	Met	Lys	Ala	Glu
			180					185					190		
Ala	Tyr	Lys	Tyr	Ile	Asp	Ile	Thr	Arg	Pro	Glu	Glu	Val	Lys	His	Gln
	195						200					205			
Glu	Gln	His	His	Pro	Glu	Leu	Glu	Asp	Lys	Asp	Leu	Asp	Phe	Lys	Asn
	210					215					220				
Lys	Asp	Thr	Asp	Ile	Ser	Ile	Lys	Pro	Glu	Gly	Val	Arg	Glu	Pro	Asp
225					230					235					240
Lys	Pro	Ala	Pro	Val	Glu	Gly	Lys	Ile	Ile	Lys	Asp	His	Leu	Leu	Glu
				245					250					255	
Glu	Ser	Thr	Phe	Ala	Pro	Tyr	Ile	Asp	Asp	Leu	Ser	Glu	Glu	Gln	Arg
			260					265					270		
Arg	Ala	Pro	Gln	Ile	Thr	Thr	Pro	Val	Lys	Ile	Thr	Leu	Thr	Glu	Ile
	275						280					285			
Glu	Pro	Ser	Val	Glu	Thr	Thr	Gln	Glu	Lys	Thr	Pro	Glu	Lys	Gln	
	290					295				300					
Asp	Ile	Cys	Leu	Lys	Pro	Ser	Pro	Asp	Thr	Val	Pro	Thr	Val	Thr	Val
305					310					315					320
Ser	Glu	Pro	Glu	Asp	Asp	Ser	Pro	Gly	Ser	Ile	Thr	Pro	Pro	Ser	Ser
				325					330					335	
Gly	Thr	Glu	Pro	Ser	Ala	Ala	Glu	Ser	Gln	Gly	Lys	Gly	Ser	Ile	Ser
			340					345					350		
Glu	Asp	Glu	Leu	Ile	Thr	Ala	Ile	Lys	Glu	Ala	Lys	Gly	Leu	Ser	Tyr
	355						360					365			
Glu	Thr	Ala	Glu	Asn	Pro	Arg	Pro	Val	Gly	Gln	Leu	Ala	Asp	Arg	Pro
	370					375					380				
Glu	Val	Lys	Ala	Arg	Ser	Gly	Pro	Pro	Thr	Ile	Pro	Ser	Pro	Leu	Asp
385					390					395					400
His	Glu	Ala	Ser	Ser	Ala	Glu	Ser	Gly	Asp	Ser	Glu	Ile	Glu	Leu	Val
				405					410					415	
Ser	Glu	Asp	Pro	Met	Ala	Ala	Glu	Asp	Ala	Leu	Pro	Ser	Gly	Tyr	Val
			420					425					430		
Ser	Phe	Gly	His	Val	Gly	Gly	Pro	Pro	Pro	Ser	Pro	Ala	Ser	Pro	Ser
	435						440					445			
Ile	Gln	Tyr	Ser	Ile	Leu	Arg	Glu	Glu	Arg	Glu	Ala	Glu	Leu	Asp	Ser
	450					455					460				
Glu	Leu	Ile	Ile	Glu	Ser	Cys	Asp	Ala	Ser	Ser	Ala	Ser	Glu	Glu	Ser
465					470					475					480
Pro	Lys	Arg	Glu	Gln	Asp	Ser	Pro	Pro	Met	Lys	Pro	Ser	Ala	Leu	Asp
				485					490					495	
Ala	Ile	Arg	Glu	Glu	Thr	Gly	Val	Arg	Ala	Glu	Glu	Arg	Ala	Pro	Ser
			500					505					510		
Arg	Arg	Gly	Leu	Ala	Glu	Pro	Gly	Ser	Phe	Leu	Asp	Tyr	Pro	Ser	Thr
	515						520					525			
Glu	Pro	Gln	Pro	Gly	Pro	Glu	Leu	Pro	Pro	Gly	Asp	Gly	Ala	Leu	Glu
	530					535					540				
Pro	Glu	Thr	Pro	Met	Leu	Pro	Arg	Lys	Pro	Glu	Glu	Asp	Ser	Ser	Ser
545					550					555					560
Asn	Gln	Ser	Pro	Ala	Ala	Thr	Lys	Gly	Pro	Gly	Pro	Leu	Gly	Pro	Gly
				565					570					575	
Ala	Pro	Pro	Pro	Leu	Leu	Phe	Leu	Asn	Lys	Gln	Lys	Ala	Ile	Asp	Leu
			580					585					590		

```

Leu Tyr Trp Arg Asp Ile Lys Gln Thr Gly Ile Val Phe Gly Ser Phe
    595                600                605
Leu Leu Leu Leu Phe Ser Leu Thr Gln Phe Ser Val Val Ser Val Val
    610                615                620
Ala Tyr Leu Ala Leu Ala Leu Ser Ala Thr Ile Ser Phe Arg Ile
    625                630                635                640
Tyr Lys Ser Val Leu Gln Ala Val Gln Lys Thr Asp Glu Gly His Pro
    645                650                655
Phe Lys Ala Tyr Leu Glu Leu Glu Ile Thr Leu Ser Gln Glu Gln Ile
    660                665                670
Gln Lys Tyr Thr Asp Cys Leu Gln Phe Tyr Val Asn Ser Thr Leu Lys
    675                680                685
Glu Leu Arg Arg Leu Phe Leu Val Gln Asp Leu Val Asp Ser Leu Lys
    690                695                700
Phe Ala Val Leu Met Trp Leu Leu Thr Tyr Val Gly Ala Leu Phe Asn
    705                710                715                720
Gly Leu Thr Leu Leu Met Ala Val Val Ser Met Phe Thr Leu Pro
    725                730                735
Val Val Tyr Val Lys His Gln Ala Gln Ile Asp Gln Tyr Leu Gly Leu
    740                745                750
Val Arg Thr His Ile Asn Ala Val Val Ala Lys Ile Gln Ala Lys Ile
    755                760                765
Pro Gly Ala Lys Arg His Ala Glu
    770                775

```

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 356 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(vii) IMMEDIATE SOURCE:

- (A) LIBRARY: GenBank
- (B) CLONE: 307309

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

```

Met Ala Ala Glu Asp Ala Leu Pro Ser Gly Tyr Val Ser Phe Gly His
 1          5          10          15
Val Gly Gly Pro Pro Pro Ser Pro Ala Ser Pro Ser Ile Gln Tyr Ser
          20          25          30
Ile Leu Arg Glu Glu Arg Glu Ala Glu Leu Asp Ser Glu Leu Ile Ile
          35          40          45
Glu Ser Cys Asp Ala Ser Ser Ala Ser Glu Glu Ser Pro Lys Arg Glu
          50          55          60
Gln Asp Ser Pro Pro Met Lys Pro Ser Ala Leu Asp Ala Ile Arg Glu
          65          70          75          80
Glu Thr Gly Val Arg Ala Glu Glu Arg Ala Pro Ser Arg Arg Gly Leu
          85          90          95
Ala Glu Pro Gly Ser Phe Leu Asp Tyr Pro Ser Thr Glu Pro Gln Pro
          100          105          110
Gly Pro Glu Leu Pro Pro Gly Asp Gly Ala Leu Glu Pro Glu Thr Pro
          115          120          125
Met Leu Pro Arg Lys Pro Glu Asp Ser Ser Ser Asn Gln Ser Pro
          130          135          140
Ala Ala Thr Lys Gly Pro Gly Pro Leu Gly Pro Gly Ala Pro Pro Pro
          145          150          155          160
Leu Leu Phe Leu Asn Lys Gln Lys Ala Ile Asp Leu Leu Tyr Trp Arg

```

[illegible]

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 208 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(vii) IMMEDIATE SOURCE:

- (A) LIBRARY: GenBank
(B) CLONE: 307311

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

Met 1	Gln	Ala	Thr	Ala 5	Asp	Ser	Thr	Lys	Met 10	Asp	Cys	Val	Trp	Ser 15	Asn
Trp	Lys	Ser	Gln 20	Ala	Ile	Asp	Leu	Leu 25	Tyr	Trp	Arg	Asp 30	Ile	Lys	Gln
Thr	Gly	Ile 35	Val	Phe	Gly	Ser	Phe 40	Leu	Leu	Leu	Leu 45	Phe	Ser	Leu	Thr
Gln	Phe 50	Ser	Val	Val	Ser	Val 55	Val	Ala	Tyr	Leu	Ala 60	Leu	Ala	Ala	Leu
Ser 65	Ala	Thr	Ile	Ser	Phe 70	Arg	Ile	Tyr	Lys	Ser 75	Val	Leu	Gln	Ala 80	Val
Gln	Lys	Thr	Asp 85	Glu	Gly	His	Pro	Phe	Lys 90	Ala	Tyr	Leu	Glu 95	Leu	Glu
Ile	Thr	Leu	Ser 100	Gln	Glu	Gln	Ile	Gln 105	Lys	Tyr	Thr	Asp 110	Cys	Leu	Gln
Phe	Tyr	Val 115	Asn	Ser	Thr	Leu	Lys 120	Glu	Leu	Arg	Arg	Leu 125	Phe	Leu	Val
Gln	Asp 130	Leu	Val	Asp	Ser	Leu 135	Lys	Phe	Ala	Val	Leu 140	Met	Trp	Leu	Leu
Thr 145	Tyr	Val	Gly	Ala	Leu 150	Phe	Asn	Gly	Leu	Thr 155	Leu	Leu	Leu	Met	Ala
Val	Val	Ser	Met	Phe	Thr	Leu	Pro	Val	Val	Tyr	Val	Lys	His	Gln	Ala

			165					170					175			
Gln	Ile	Asp	Gln	Tyr	Leu	Gly	Leu	Val	Arg	Thr	His	Ile	Asn	Ala	Val	
			180					185					190			
Val	Ala	Lys	Ile	Gln	Ala	Lys	Ile	Pro	Gly	Ala	Lys	Arg	His	Ala	Glu	
		195					200					205				

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 267 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(vii) IMMEDIATE SOURCE:

- (A) LIBRARY: GenBank
(B) CLONE: 281046

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

Met 1	Asp	Cys	Val	Trp 5	Ser	Asn	Trp	Lys	Ser 10	Gln	Ala	Ile	Asp	Leu 15	Leu
Tyr	Trp	Arg	Asp 20	Ile	Lys	Gln	Thr	Gly 25	Ile	Val	Phe	Gly	Ser 30	Phe	Leu
Leu	Leu	Leu 35	Phe	Ser	Leu	Thr	Gln 40	Phe	Ser	Val	Val	Ser 45	Val	Val	Ala
Tyr	Leu 50	Ala	Leu	Ala	Ala	Leu 55	Ser	Ala	Thr	Ile	Ser 60	Phe	Arg	Ile	Tyr
Lys 65	Ser	Val	Leu	Gln 70	Ala	Val	Gln	Lys	Thr	Asp 75	Glu	Gly	His	Pro	Phe 80
Lys	Ala	Tyr	Leu	Glu 85	Leu	Glu	Ile	Thr	Leu 90	Ser	Gln	Glu	Gln	Ile 95	Gln
Lys	Tyr	Thr	Asp 100	Cys	Leu	Gln	Leu	Tyr 105	Val	Asn	Ser	Thr	Leu 110	Lys	Glu
Leu	Arg	Arg 115	Leu	Phe	Leu	Val	Gln 120	Asp	Leu	Val	Asp	Ser 125	Leu	Lys	Phe
Ala	Val 130	Leu	Met	Trp	Leu	Leu 135	Thr	Tyr	Val	Gly	Ala 140	Leu	Phe	Asn	Gly
Leu 145	Thr	Leu	Leu	Leu 150	Met	Ala	Val	Val	Ser	Met 155	Phe	Thr	Leu	Pro	Val 160
Val	Tyr	Val	Lys 165	His	Gln	Ala	Gln	Val	Asp 170	Gln	Tyr	Leu	Gly	Leu 175	Val
Arg	Thr	His 180	Ile	Asn	Thr	Val	Val	Ala 185	Lys	Ile	Gln	Ala	Lys 190	Ile	Pro
Gly	Ala	Arg 195	Gly	Met	Leu	Ser	Arg 200	Trp	Leu	Pro	Gln	Glu 205	Lys	Pro	Asp
Met	Asn 210	Gly	Gly	Val	Trp	Ser 215	Gly	Asn	Ser	Ser	Leu 220	Leu	Pro	Arg	Tyr
Cys 225	Glu	Leu	Ile	Val 230	Ser	Leu	Pro	Gln	Tyr	His 235	Asn	Leu	Arg	Gly	Lys 240
Leu	Arg	Asp	Arg 245	Cys	Phe	Gln	Ser	Phe	Pro 250	Val	Leu	Leu	Gly	Tyr 255	Leu
Ser	Pro	Pro	Arg 260	Pro	Leu	Ser	Ser	Thr 265	Lys	Val					

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 261 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vii) IMMEDIATE SOURCE:

(A) LIBRARY: SPLNFET01

(B) CLONE: 28742

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

CCTATNCCNG	CTGCTTTCAT	TGACAGTATT	CAGCATTGTG	AGCGTAACAG	CCTACATTGC	60
CTTNGCCCTG	CNCTCTGTGA	CCATCAGCTN	TAGGCTATAC	AAGGGTGTGA	TCCAAGCTAT	120
CCAGAAATCA	GATGAAGGNC	ACCCATTTCAG	GGCATATCTG	GANTCTGAAG	TTGCTATATC	180
TGAGGAGTTG	NTTCAGAAGT	ACACGTAAAT	NNTGNNCNTG	GTCAATGTGA	NCTCCACGNC	240
TAANGGANCT	CAGGTGCCTA	T				261

CLAIMS

1. A substantially purified human NSPLP protein comprising the amino acid sequence of SEQ ID NO:1 or fragments thereof.

2. An isolated and purified polynucleotide sequence encoding a protein of claim 1.

3. An isolated and purified polynucleotide sequence of claim 2 consisting of the sequence of SEQ ID NO:2 or degenerate variants thereof.

4. A polynucleotide sequence fully complementary to the sequence of SEQ ID NO:2 or degenerate variants thereof.

5. An isolated and purified polynucleotide sequence of claim 2 consisting of a polynucleotide sequence that hybridizes under stringent hybridization conditions to the sequence of SEQ ID NO:2.

6. A recombinant expression vector containing a polynucleotide sequence of claim 2.

7. A recombinant host cell comprising a polynucleotide sequence of claim 2.

8. A method for producing a polypeptide comprising the amino acid sequence shown in SEQ ID NO:1, the method comprising the steps of:

a) culturing the host cell of Claim 7 under conditions suitable for the expression of the polypeptide; and

b) recovering the polypeptide from the host cell culture.

9. A recombinant expression vector containing a polynucleotide sequence of claim 5.

10. A recombinant host cell comprising a polynucleotide sequence of claim 9.

11. A pharmaceutical composition comprising a substantially purified human NSPLP protein (SEQ ID NO:1) in conjunction with a suitable pharmaceutical carrier.

12. A purified antibody which binds specifically to a polypeptide of claim 1.

13. A purified antagonist which specifically blocks or reduces the activity of the polypeptide of claim 1.

14. A pharmaceutical composition comprising a substantially purified antagonist of the polypeptide of claim 1 in conjunction with a suitable pharmaceutical carrier.

15. A substantially purified human NSPLP protein comprising the amino acid sequence of SEQ ID NO:3 or fragments thereof.

16. An isolated and purified polynucleotide sequence encoding a protein of claim 15.

17. An isolated and purified polynucleotide sequence of claim 16 consisting of the sequence of SEQ ID NO:4 or degenerate variants thereof.

18. A polynucleotide sequence fully complementary to the sequence of SEQ ID NO:4 or degenerate variants thereof.

19. An isolated and purified polynucleotide sequence of claim 16 consisting of a polynucleotide sequence that hybridizes under stringent hybridization conditions to the sequence of SEQ ID NO:4.

20. A recombinant expression vector containing a polynucleotide sequence of claim 16.

21. A recombinant host cell comprising a polynucleotide sequence of claim 16.

22. A method for producing a polypeptide comprising the amino acid sequence shown in SEQ ID NO:3, the method comprising the steps of:

a) culturing the host cell of Claim 21 under conditions suitable for the expression of the polypeptide; and

b) recovering the polypeptide from the host cell culture.

23. A recombinant expression vector containing a polynucleotide sequence of claim 19.

24. A recombinant host cell comprising a polynucleotide sequence of claim 23.

25. A pharmaceutical composition comprising a substantially purified human NSPLP protein (SEQ ID NO:3) in conjunction with a suitable pharmaceutical carrier.

26. A purified antibody which binds specifically to a polypeptide of claim 15.

27. A purified antagonist which specifically blocks or reduces the activity of the polypeptide of claim 15.

28. A pharmaceutical composition comprising a substantially purified antagonist of the polypeptide of claim 15 in conjunction with a suitable pharmaceutical carrier.

1/24

11 5' TTT GTG CAG TTA CAG CTT TTC TTT TGG TAT GCA TAA TTA ATA NTT GGA GCT GCA 56
 47
 65 65 AAG AGA TCG TGA CAA GAG ATG GAC GGT CAG AAG AAA AAT TGG AAG GAC AAG GTT 110
 74 83 M D G Q K K N W K D K V
 119 119 GTT GAC CTC CTG TAC TGG AGA GAC ATT AAG AAG ACT GGA GTG GTG TTT GGT GCC 164
 128 137 V D L L Y W R D I K K T G V V F G A
 146 155
 173 173 AGC CTA TTC CTG CTG CTT TCA TTG ACA GTA TTC AGC ATT GTG AGC GTA ACA GCC 218
 182 191 S L F L L L S L T V F S I V S V T A
 227 227 TAC ATT GCC TTG GCC CTG CTC TCT GTG ACC ATC AGC TTT AGG ATA TAC AAG GGT 272
 236 245 Y I A L A L L S V T I S F R I Y K G
 254 263
 281 281 GTG ATC CAA GCT ATC CAG AAA TCA GAT GAA GGC CAC CCA TTC AGG GCA TAT CTG 326
 290 299 V I Q A I Q K S D E G H P F R A Y L
 308 317
 335 335 GAA TCT GAA GTT GCT ATA TCT GAG GAG TTG GTT CAG AAG TAC AGT AAT TCT GCT 380
 344 353 E S E V A I S E E L V Q K Y S N S A
 362 371

Fig. 1A

2/24

389	398	407	416	425	432
CTT GGT CAT GTG AAC TGC ACG ATA AAG GAA CTC AGG CGC CTC TTC TTA GTT GAT					
L G H V N C T I K E L R R L F L V D					
443	452	461	470	479	488
GAT TTA GTT GAT TCT CTG AAG TTT GCA GTG TTG ATG TGG GTA TTT ACC TAT GTT					
D L V D S L K F A V L M W V F T Y V					
497	506	515	524	533	542
GGT GCC TTG TTT AAT GGT CTG ACA CTA CTG ATT TTG GCT CTC ATT TCA CTC TTC					
G A L F N G L T L L I L A L I S L F					
551	560	569	578	587	596
AGT GTT CCT GTT ATT TAT GAA CGG CAT CAG GCA CAG ATA GAT CAT TAT CTA GGA					
S V P V I Y E R H Q A Q I D H Y L G					
605	614	623	632	641	650
CTT GCA AAT AAG AAT GTT AAA GAT GCT ATG GCT AAA ATC CAA GCA AAA ATC CCT					
L A N K N V K D A M A K I Q A K I P					
659	668	677	686	695	704
GGA TTG AAG CGC AAA GCT GAA TGA AAA CGC CCA AAA TAA TTA GTA GGA GTT CAT					
G L K R K A E					

Fig. 1B

3/24

713 722 731 740 749 758
CTT TAA AGG GGA TAT TCA TTT GAT TAT ACG GGG GAG GGT CAG GGA AGA ACG ACC

767 776 785 794
TTG ACG TTG CAG TGC AGT TTC ACA GAT CGT TGT TAG ATC TT 3'

Fig. 1C

4/24

9	18	27	36	45	54
5' CAC NAG CGN NTC GNG CTC CCG AAC CTC TAG CTG CGA CTC GGA NTG AGT CAG TCA					
63	72	81	90	99	108
GTC TGT CGG AGT CTG TCC TCG GAG CAG GCG GAG TAA AGG GAC TTG AGC GAG CCA					
117	126	135	144	153	162
GTT GCC GGA TTA TTC TAT TTC CCC TCC CTC TCT CCC GCC CCG TAT CTC TTT TCA					
171	180	189	198	207	216
TTT TNN TNC CAC CCT TGC TCG CGT ANC ATG GCG GAG CGT NCG GCG GCC ACT CAG					
		M	A	E	R
			A	X	A
					T
					Q
225	234	243	252	261	270
TCC CAT TCC ATC TCC TCG TCG TCC TCC GGA GCC GAG CCG TCC GCG CCC GGC GGC					
S H S I S S S S F G A E P S A P G G					
279	288	297	306	315	324
GGC GGC AGC CCA GGA GCC TGC CCC GCG CTG GCG GCG AAG AGC TGC AGC TCC TCC					
G G S P G A C P A L G T K S C S S					
333	342	351	360	369	378
TGT GCG GTG CAC GAT CTG ATT TTM TGG AGA GAT GTG AAG AAG ACT GGC TTT GTC					
C A V H D L I X W R D V K K T G F V					

Fig. 2A

5/24

387	396	405	414	423	432
TTT GGC ACC	ACG CTG ATC	ATG CTG CTT	TCC CTG GCA	GCT TTC AGT	GTC ATC AGT
F G T	T L I	M L L	S L A	F S V	I S
441	450	459	468	477	486
GTG GTT TCT	TAC CTC ATC	CTG GCT CTT	CTC TCT GTC	ACC ATC AGC	TTC AGG ATC
V V S	Y L I	L A L	S V T	I S F	R I
495	504	513	522	531	540
TAC AAG TCC	GTC ATC CAA	GCT GTA CAG	AAG TCA GAA	GAA GGC CAT	CCA TTC AAA
Y K S	V I Q	A V Q	K S E	G H P	F K
549	558	567	576	585	594
GCC TAC CTG	GAC GTA	ATT ACT	CTG TCC	TCA GAA	GCT TTC
A Y L	D V D	I T L	S S E	A F H	N Y M
603	612	621	630	639	648
AAT GCT GCC	ATG GTG	CAC ATC	AAC AGG	GCC CTG	AAA CTC
N A A	M V H	I N R	A L K	L I I	R L F
657	666	675	684	693	702
CTG GTA GAA	GAT CTG	GTT GAC	TCC TTG	AAG CTG	GCT GTC
L V E	D L V	D S L	K L A	V F M	W L M
711	720	729	738	747	756
ACC TAT GTT	GGT GCT	GTT TTT	AAC GGA	ATC ACC	CTT CTA
T Y V	G A V	F N G	I T L	L I L	A A E

Fig. 2B

6/24

765	774	783	792	801	810
CTC ATT TTN AGT GTC CCG ATT GTN TAT NAG AAG TAC AAG GTT CCA AGC AAA ACT					
L I X S V P I V Y X K Y K V P S K T					
819	828	837	846	855	864
CCC TGG AAT CGC CAA AAA AAA GGC AGA ATA AGT ACA TGG AAA CCA GAA ATG CAA					
P W N R Q K K K G R I S T W K P E M Q					
873	882	891	900	909	918
CAG TTA CTA AAA CAC CAT TTA ATA GTT ATA ACG TCG TTA CTT GTA CTA TGA AGG					
Q L L K H H L I V I T S L L V L					
927	936	945	954	963	972
AAA ATA CTC AGT GTC AGC TTG AGC CTG CAT TCC AAG CTT TTT TAA TTT GGT					
981	990	999	1008	1017	1026
GGT TTT CTC CCA TCC TTT CCC TTT AAC CCT CAG TAT CAA GCA CAA ANT TTN ATG					
1035	1044	1053	1062	1071	1080
GAC TGA TAA NNG ATC TAT NTT AGA NCT CAG AAG ANG ANA GNT TCA NNT GCA TAG					
1089					
GNT AAG GNA NTA CC 3'					

Fig. 2C

7/24

CORNNOT01	corneal fibroblasts, 76y	4	0.3996
FIBRAGT02	ATGD60 fibroblasts, ataxia telan, radiation 30 min	2	0.3968
BLADNOT01	bladder, 78 F	10	0.3494
OVARNON01	ovary, 59 F, NORM	2	0.3185
U937NOT01	U937 monocyte cell line, 37 M	6	0.2973
FIBRNOT01	WI38 lung fibroblast cell line, fetal F	6	0.2812
SCORNON02	spinal cord, 71 M, NORM	7	0.2415
COCHFEM01	ear, cochlea, fetal, WM	2	0.2315
KIDNNOT01	kidney, 64 F	1	0.1562
BRSTTUT02	breast tumor, 54 F, match to BRSTNOT03	5	0.1489
PANCTUT02	pancreatic tumor, carcinoma, 45 F	5	0.1402
SCORNON01	spinal cord, 71 M, NORM	1	0.1379
BRSTNOT01	breast, 56 F	7	0.1345
BRAINOT03	brain, 26 M	7	0.1297
BLADTUT02	bladder tumor, carcinoma, 80 F	4	0.1220
PROSTUT05	prostate tumor, 69 M, match to PROSNOT07	4	0.1212
BSTMNOT01	brain stem, 72 M	1	0.1203
PROSTUT04	prostate tumor, 57 M, match to PROSNOT06	7	0.1140
MPHGNOT03	macrophages (adher PBMNC), M/F	8	0.1032
HMC1NOT01	HMC-1 mast cell line, 52 F	3	0.0999
SPLNFEM01	spleen, fetal, WM	3	0.0995
MMLR3DT01	macrophages (adher PBMNC), M/F, 72-hr MLR	3	0.0993
BRAINOT11	brain, right temporal, epilepsy, 5 M	3	0.0966
MMLR1DT01	macrophages (adher PBMNC), M/F, 24-hr MLR	4	0.0944
NEUTGMT01	granulocytes, periph blood, M/F, treated GM-CSF	6	0.0938
RETNNOM02	retina, 55 M, NORM, WM	1	0.0899
MUSCNOT01	muscle, skeletal	2	0.0888
RATRNOT01	heart, right atrium, 51 F	1	0.0861

Fig. 3A

8/24

LUNGNOM01	lung, 72 M, WM	3	0.0802
BRAINOT09	brain, fetal M	3	0.0783
MUSCNOT02	muscle, psoas, 12 M	2	0.0763
TLYMNOT02	lymphocytes (non-adher PBMNC), M/F	3	0.0761
PROSNOT01	prostate, 78 M	2	0.0696
NEUTLPT01	granulocytes, periph blood, M/F, treated LPS	4	0.0671
UTRSNOT02	uterus, 34 F	4	0.0666
BSTMNON02	brain stem, 72 M, NORM	2	0.0637
STOMNOT02	stomach, 52 M, match to STOMTUT01	2	0.0615
SCORNOT01	spinal cord, 71 M	3	0.0603
TONGTUT01	tongue tumor, carcinoma, 36 M	2	0.0590
BRAITUT08	brain tumor, astrocytoma, 47 M	4	0.0582
PROSNOT05	prostate, 67 M, match to PROSTUT03	1	0.0575
LATRTUT02	heart tumor, myoma, 43 M	4	0.0548
SYNOOAT01	synovium, knee, osteoarthritis, 82 F	3	0.0538
HIPONOT01	brain, hippocampus, 72 F	1	0.0535
PENITUT01	penis tumor, carcinoma, 64 M	2	0.0532
KIDNNOT09	kidney, fetal M	2	0.0531
SYNORAT04	synovium, wrist, rheumatoid, 62 F	3	0.0521
BRAITUT02	brain tumor, metastasis, 58 M	3	0.0507
MENITUT03	brain tumor, benign meningioma, 35 F	2	0.0499
PLACNOT02	placenta, fetal F	3	0.0495
THP1PEB01	THP-1 promonocyte cell line, treated PMA	1	0.0487
MPHGNOT02	macrophages (adher PBMNC), 24 M	1	0.0478
BRSTNOT02	breast, 55 F, match to BRSTTUT01	3	0.0475
RATRNOT02	heart, right atrium, 39 M	2	0.0472
COLNCRT01	colon, Crohn's, 40 M, match to COLNNOT05	1	0.0468

Fig. 3B

9/24

LVENNOT01	heart, left ventricle, 51 F	1	0.0450
PLACNOB01	placenta, neonatal F	2	0.0450
BRSTTUT03	breast tumor, 58 F, match to BRSTNOT05	3	0.0444
PLACNOM02	placenta, neonatal F, NORM, WM	8	0.0444
COLNNOT01	colon, 75 M, match to COLNTUT02	2	0.0426
BRSTNOM02	breast, F, NORM, WM	2	0.0413
THP1PLB02	THP-1 promonocyte cell line, treated PMA, LPS	1	0.0407
BRAITUT01	brain tumor, oligoastrocytoma, 50 F	3	0.0403
KERANOT02	keratinocytes, primary cell line, 30 F	1	0.0396
THYMNOT02	thymus, 3 M	2	0.0386
HNT2AGT01	hNT-2 cell line, post-mitotic neurons	2	0.0380
STOMTUT01	stomach tumor, 52 M, match to STOMNOT02	1	0.0367
MMLR2DT01	macrophages (adher PBMNC), M/F, 48-hr MLR	2	0.0354
SPLNFET01	spleen, fetal	1	0.0352
PROSNOT07	prostate, 69 M, match to PROSTUT05	1	0.0347
TONSNOT01	tonsil, hyperplasia, 6 M	1	0.0339
LUNGNOT01	lung, 72 M	1	0.0338
PROSNOT06	prostate, 57 M, match to PROSTUT04	1	0.0332
PGANNOT03	paraganglia, 46 M	1	0.0309
PROSTUT01	prostate tumor, 50 M, match to PROSNOT02	1	0.0309
BRAITUT03	brain tumor, astrocytoma, 17 F	2	0.0307
BEPINOT01	bronchial epithelium, primary cell line, 54 M	1	0.0304
BRSTTUT01	breast tumor, 55 F, match to BRSTNOT02	2	0.0302
STOMNOT01	stomach, 55 M	1	0.0300
BRAINOT12	brain, right frontal, epilepsy, 5 M	1	0.0299
BRSTNOT03	breast, 54 F, match to BRSTTUT02	2	0.0293
SYNORAT05	synovium, knee, rheumatoid, 62 F	1	0.0286

Fig. 3C

10/24

LUNGNOT09	lung, fetal M	1	0.0285
TESTTUT02	testicular tumor, 31 M	1	0.0278
THYRNOT03	thyroid tumor, adenoma, 28 F	2	0.0276
COLNTUT03	colon tumor, 62 M, match to COLNNOT16	1	0.0272
HYPONOB01	hypothalamus, 16-75 M/F	1	0.0270
BRSTNOM01	breast, F, NORM, WM	1	0.0264
LATRNOT01	heart, left atrium, 51 F	1	0.0263
LIVRNOM01	liver, 49 M, WM	1	0.0254
PANCNOT08	pancreas, 65 F, match to PANCTUT01	1	0.0254
TMLR3DT01	lymphocytes (non-adher PBMNC), M, 96-hr MLR	1	0.0229
SPLNNOT02	spleen, 29 M	1	0.0220
MELANOM01	melanocytes, M, NORM, WM	2	0.0216
TMLR2DT01	lymphocytes (non-adher PBMNC), M/F, 24-hr MLR	1	0.0211
EOSIHET02	eosinophils, hypereosinophilia, 48 M	2	0.0209
LUNGNOT03	lung, 79 M, match to LUNGTUT02	1	0.0200
SYNORAB01	synovium, hip, rheumatoid, 68 F	1	0.0195
CERVNOT01	cervix, 35 F	1	0.0193
LUNGTUT02	lung tumor, metastasis, 79 M, match to LUNGNOT03	1	0.0189
LUNGNOT04	lung, 2 M	1	0.0183
NEUTFMT01	granulocytes, periph blood, M/F, treated fMLP	1	0.0174
KIDNNOT05	kidney, neonatal F	1	0.0161
PGANNOT01	paraganglia, 46 M	1	0.0160
NGANNOT01	ganglioneuroma, 9 M	1	0.0155
BRSTNOT05	breast, 58 F, match to BRSTTUT03	1	0.0154
CORPNOT02	brain, corpus callosum, Alzheimer's, 74 M	1	0.0151
COLNFET02	colon, fetal F	1	0.0142
LUNGFET03	lung, fetal F	1	0.0137

Fig. 3D

11/24

UCMCL5T01	lymphocytes (umbilical cord), treated IL-5	1	0.0125
LIVSFEM02	liver/spleen, fetal M, NORM, WM	3	0.0087
BRAINOM01	brain, infant F, NORM, WM	1	0.0045

Fig. 3E

12/24

ADRENOT01	adrenal gland, 10-46 M/F	2	0.2081
BRAINOT03	brain, 26 M	11	0.2039
NEUTLPT01	granulocytes, periph blood, M/F, treated LPS	10	0.1678
SCORNON01	spinal cord, 71 M, NORM	1	0.1379
OLFENOM01	epithelium, olfactory, 35 F, WM	1	0.1330
THP1NOB01	THP-1 promonocyte cell line, control	4	0.1309
BMARNOR02	bone marrow, 16-70 M/F, RP	3	0.1294
BRAINOT11	brain, right temporal, epilepsy, 5 M	4	0.1288
BRSTNOM02	breast, F, NORM, WM	6	0.1239
HIPONOT01	brain, hippocampus, 72 F	2	0.1070
BRAINOT04	brain, choroid plexus, hemorrhage, 44 M	3	0.1067
SCORNON02	spinal cord, 71 M, NORM	3	0.1035
LUNGNOT01	lung, 72 M	3	0.1014
THP1PEB01	THP-1 promonocyte cell line, treated PMA	2	0.0975
KIDNNOT02	kidney, 64 F	2	0.0964
BRSTNOT01	breast, 56 F	5	0.0960
PITUNOT02	pituitary, 7-65 M/F	2	0.0905
RETNNOM02	retina, 55 M, NORM, WM	1	0.0899
BRAINOT12	brain, right frontal, epilepsy, 5 M	3	0.0898
MUSCNOT02	muscle, psoas, 12 M	2	0.0763
OVARNOM01	ovary, 49 F, WM	1	0.0752
EOSIHET02	eosinophils, hypereosinophilia, 48 M	7	0.0731
HEARNOT01	heart, 56 M	1	0.0707
KIDNNOT05	kidney, neonatal F	4	0.0645
NEUTGMT01	granulocytes, periph blood, M/F, treated GM-CSF	4	0.0625
BRSTNOT07	breast, 43 F	2	0.0614

Fig. 4A

13/24


SCORNOT01	spinal cord, 71 M	3	0.0603
HNT2RAT01	hNT-2 cell line, teratocarcinoma, treated RA	3	0.0556
LARTTUT02	heart tumor, myoma, 43 M	4	0.0548
HUVELPB01	HUVEC endothelial cell line, treated cytokine, LPS	1	0.0546
BRAITUT01	brain tumor, oligoastrocytoma, 50 F	4	0.0537
PANCNOT04	pancreas, 5 M	1	0.0504
PLACNOT02	placenta, fetal F	3	0.0495
RATRN0T02	heart, right atrium, 39 M	2	0.0472
BRAINOM02	brain, 55 M, NORM, WM	1	0.0454
MELANOM01	melanocytes, M, NORM, WM	4	0.0431
HUVENOB01	HUVEC endothelial cell line, control	1	0.0419
THYRN0T03	thyroid tumor, adenoma, 28 F	1	0.0386
OVARNOT03	ovary, 43 F, match to OVARTUT01	1	0.0386
THYMN0T02	thymus, 3 M	2	0.0386
HNT2AGT01	hNT-2 cell line, post-mitotic neurons	2	0.0380
PROSN0T07	prostate, 69 M, match to PROSTUT05	1	0.0347
SYNORAT03	synovium, wrist, rheumatoid, 56 F	2	0.0339
LNODNOT02	lymph nodes, 42 F	1	0.0335
TBLYNOT01	T-B lymphoblast cell line, leukemia	1	0.0326
OVARTUT01	ovarian tumor, 43 F, match to OVARNOT03	1	0.0323
BSTMNON02	brain stem, 72 M, NORM	1	0.0319
OVARNOT02	ovary, 59 F	1	0.0315
BLADTUT02	bladder tumor, carcinoma, 80 F	1	0.0305
LUNGTUT03	lung tumor, carcinoma, 69 M	1	0.0305
CORPN0T02	brain, corpus callosum, Alzheimer's, 74 M	2	0.0302
BRSTTUT02	breast tumor, 54 F, match to BRSTNOT03	1	0.0298

Fig. 4B

14/24

SYNORAT05	synovium, knee, rheumatoid, 62 F	1	0.0286
LUNGNOT09	lung, fetal M	1	0.0285
COLNFET02	colon, fetal F	2	0.0284
PLACNOM02	placenta, neonatal F, NORM, WM	5	0.0278
LUNGNOT12	lung, 78 M	1	0.0277
BRAINOT09	brain, fetal M	1	0.0261
LUNGNOT10	lung, fetal M	1	0.0261
BRAITUT07	brain tumor, left frontal, 32 M	1	0.0258
TLYMNOT02	lymphocytes (non-adher PBMNC), M/F	1	0.0254
LUNGNOT02	lung, 47 M	1	0.0245
MLLR1DT01	macrophages (adher PBMNC), M/F, 24-hr MLR	1	0.0236
TMLR3DT01	lymphocytes (non-adher PBMNC), M, 96-hr MLR	1	0.0229
PLACNOB01	placenta, neonatal F	1	0.0225
CRBLNOT01	brain, cerebellum, 69 M	1	0.0195
CERVNOT01	cervix, 35 F	1	0.0193
ADENINB01	adenoid, inflamed, 3y	1	0.0190
LUNGNOT02	lung tumor, metastasis, 79 M, match to LUNGNOT03	1	0.0189
SYNOOAT01	synovium, knee, osteoarthritis, 82 F	1	0.0179
NEUTFMT01	granulocytes, periph blood, M/F, treated fMLP	1	0.0174
UTRSNOT02	uterus, 34 F	1	0.0166
PGANNOT01	paraganglia, 46 M	1	0.0160
BRAITUT03	brain tumor, astrocytoma, 17 F	1	0.0153
BRSTTUT03	breast tumor, 58 F, match to BRSTNOT05	1	0.0148
BRSTNOT03	breast, 54 F, match to BRSTTUT02	1	0.0147
LUNGFET03	lung, fetal F	1	0.0137
LIVSFEM02	liver/spleen, fetal M, NORM, WM	4	0.0116
BRAINOM01	brain, infant F, NORM, WM	2	0.0089

Fig. 4C



7 Fragments in 1 Contigs

Fig. 5

16/24

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100	101	102	103	104	105	106	107	108	109	110	111	112	113	114	115	116	117	118	119	120	121	122	123	124	125	126	127	128	129	130	131	132	133	134	135	136	137	138	139	140	141	142	143	144	145	146	147	148	149	150	151	152	153	154	155	156	157	158	159	160	161	162	163	164	165	166	167	168	169	170	171	172	173	174	175	176	177	178	179	180	181	182	183	184	185	186	187	188	189	190	191	192	193	194	195	196	197	198	199	200	201	202	203	204	205	206	207	208	209	210	211	212	213	214	215	216	217	218	219	220	221	222	223	224	225	226	227	228	229	230	231	232	233	234	235	236	237	238	239	240	241	242	243	244	245	246	247	248	249	250	251	252	253	254	255	256	257	258	259	260	261	262	263	264	265	266	267	268	269	270	271	272	273	274	275	276	277	278	279	280	281	282	283	284	285	286	287	288	289	290	291	292	293	294	295	296	297	298	299	300	301	302	303	304	305	306	307	308	309	310	311	312	313	314	315	316	317	318	319	320	321	322	323	324	325	326	327	328	329	330	331	332	333	334	335	336	337	338	339	340	341	342	343	344	345	346	347	348	349	350	351	352	353	354	355	356	357	358	359	360	361	362	363	364	365	366	367	368	369	370	371	372	373	374	375	376	377	378	379	380	381	382	383	384	385	386	387	388	389	390	391	392	393	394	395	396	397	398	399	400	401	402	403	404	405	406	407	408	409	410	411	412	413	414	415	416	417	418	419	420	421	422	423	424	425	426	427	428	429	430	431	432	433	434	435	436	437	438	439	440	441	442	443	444	445	446	447	448	449	450	451	452	453	454	455	456	457	458	459	460	461	462	463	464	465	466	467	468	469	470	471	472	473	474	475	476	477	478	479	480	481	482	483	484	485	486	487	488	489	490	491	492	493	494	495	496	497	498	499	500	501	502	503	504	505	506	507	508	509	510	511	512	513	514	515	516	517	518	519	520	521	522	523	524
---	---	---	---	---	---	---	---	---	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

Fig. 6A

17/24

[illegible]

Fig. 6B

18/24

[illegible]

Fig. 6C

64	YKGV	IQA	IQA	IK	SD	DE	GH	PF	RAY	LE	SE	VA	IS	EL	VQ	YK	SN	SA	L	SEQ ID NO-1
100	YKSV	IQA	VQK	SE	EG	HP	FK	AY	LD	VD	IT	LS	SE	AF	HN	YM	NA	AM		SEQ ID NO-3
641	YKSV	LQA	VQK	TD	EG	HP	FK	AY	LE	IE	IT	LS	QE	QI	QK	YT	DC	LQ		SEQ ID NO-5
221	YKSV	LQA	VQK	TD	EG	HP	FK	AY	LE	IE	IT	LS	QE	QI	QK	YT	DC	LQ		SEQ ID NO-6
73	YKSV	LQA	VQK	TD	EG	HP	FK	AY	LE	IE	IT	LS	QE	QI	QK	YT	DC	LQ		SEQ ID NO-7
64	YKSV	LQA	VQK	TD	EG	HP	FK	AY	LE	IE	IT	LS	QE	QI	QK	YT	DC	LQ		SEQ ID NO-8
104	GHVN	CT	IK	EL	RR	LF	LV	DD	VD	SL	KL	FA	VL	MW	VF	TY	VG	AL	FN	SEQ ID NO-1
140	VHIN	RAL	KL	II	RI	FL	VE	DD	VD	SL	KL	AV	FL	MW	LM	TY	VG	AL	FN	SEQ ID NO-3
681	FYVN	STL	KE	LR	RL	FL	VQ	DD	VD	SL	KL	FA	VL	MW	LL	TY	VG	AL	FN	SEQ ID NO-5
261	FYVN	STL	KE	LR	RL	FL	VQ	DD	VD	SL	KL	FA	VL	MW	LL	TY	VG	AL	FN	SEQ ID NO-6
113	FYVN	STL	KE	LR	RL	FL	VQ	DD	VD	SL	KL	FA	VL	MW	LL	TY	VG	AL	FN	SEQ ID NO-7
104	LYVN	STL	KE	LR	RL	FL	VQ	DD	VD	SL	KL	FA	VL	MW	LL	TY	VG	AL	FN	SEQ ID NO-8
144	GLTL	LL	IA	LS	LF	SV	PV	IY	ER	HQ	AI	DI	HY	LG	LA	AN	KN	VK	DA	SEQ ID NO-1
180	GI	TL	LL	IA	EL	IX	SV	PI	VY	XK	Y									SEQ ID NO-3
721	GLTL	LL	MA	VV	SM	FT	LP	VV	VY	VK	HQ	AI	DI	QY	LG	LV	RT	HI	NA	SEQ ID NO-5
301	GLTL	LL	MA	VV	SM	FT	LP	VV	VY	VK	HQ	AI	DI	QY	LG	LV	RT	HI	NA	SEQ ID NO-6
153	GLTL	LL	MA	VV	SM	FT	LP	VV	VY	VK	HQ	AI	DI	QY	LG	LV	RT	HI	NA	SEQ ID NO-7
144	GLTL	LL	MA	VV	SM	FT	LP	VV	VY	VK	HQ	AI	DI	QY	LG	LV	RT	HI	NT	SEQ ID NO-8
184	MA	KI	QA	KI	P															SEQ ID NO-1
202	-	-	KV	PS	KT	P	W	NR	QK	GR	IS	TW	-	-	KP	EM	QQ	-	-	SEQ ID NO-3
761	VAKI	QA	KI	P																SEQ ID NO-5
341	VAKI	QA	KI	P																SEQ ID NO-6
193	VAKI	QA	KI	P																SEQ ID NO-7
184	VAKI	QA	KI	P																SEQ ID NO-8

Fig. 6E

194	L	K	R	K	A	E
228	L	K	H	I	V	I T S L - - - - - L V L
773	-	-	R	H	A	E
353	-	-	R	H	A	E
205	-	-	R	H	A	E
221	L	P	R	Y	C	E
						L I V S L P Q Y H N L R G K L R D R C F Q S F P V L L G Y L S P P R
199						SEQ ID NO-1
241						SEQ ID NO-3
776						SEQ ID NO-5
356						SEQ ID NO-6
208						SEQ ID NO-7
261	P	L	S	S	T	K V
						SEQ ID NO-8

Fig. 6F

22/24




Fig. 7




Fig. 8




Fig. 9